

CHIMERIC POLYPEPTIDES ALLOWING EXPRESSION OF  
PLANT-NOXIOUS PROTEINS

## FIELD OF THE INVENTION

5 This invention relates to chimeric polypeptides comprising vacuole targeting sequences and plant-noxious sequences and especially pest control proteins. The polypeptides are useful in methods for targeting non-vacuolar harmful proteins to plant vacuoles. Chimeric polypeptides of the invention containing pest control proteins are useful for conferring  
10 pest resistance on plants and in the production of compositions useful as pesticides. The methods and compositions form further aspects of the invention.

## BACKGROUND OF THE INVENTION

15 Expression of proteins in plants is a useful strategy for producing commercial quantities of a desired protein. Plant expression may avoid problems associated with production of those proteins in animal systems particularly where the protein is required for human therapeutic purposes, and can also be useful for conferring beneficial properties on the plant expressing same. Such beneficial properties may include herbicide or pest resistance  
20 for example.

However, proteins desirable for expression in plants may themselves be noxious to the plant. That is, they may harm the plant by killing or damaging it or interfering with growth, development and fertility. For example, the protein avidin has been shown to cause male  
25 sterility when expressed in plants (WO 96/40949 and WO 99/04023), as has ribonuclease when used under specific promoters (Mariani *et al.*, *Symp. Soc. Exp. Biol.* 45:271-9, 1991).

Accordingly, there is a need for a means of producing desirable plant-noxious proteins in a plant. Organelle targeting of proteins has been contemplated (US 5,792,923). Targeting  
30 of foreign proteins to vacuoles has also been contemplated. Vacuole targeting has been applied to increasing accumulation in vacuoles of products which would otherwise be metabolised. US 5,436,394 discusses targeting of invertase to the vacuole as does WO 92/14832. US 5,792,923 discloses plants in which a polyfructan sucrose is targeted to vacuoles. In US 5,723,764 cellulose synthase is targeted to vacuoles. None of these products  
35 are plant-noxious. Accordingly, there is no suggestion in any of these documents that vacuole targeting is required to avoid harmful effects on plants.

US 5,360,726 and US 5,525,713 contemplate vacuolar targeting of cereal lectins in leaves and other tissues. Lectins are themselves vacuolar proteins normally located in root tips  
40 of adult plants, and specific cells of developing embryos. Lectins are insecticidal proteins. However, there is no suggestion in any of these US patents that vacuolar targeting is necessary

or advantageous for production of insecticidal plants.

5 In WO 98/11235 it is suggested that cellulose degrading enzymes be targeted to vacuoles of transgenic plants to alleviate toxicity problems. However, no data is presented on cellulase activity or localisation of the protein in transgenic plants. Accordingly, there is no data showing vacuolar accumulation occurred and that toxicity was avoided. Therefore, there is still a need for production of transgenic plants in which plant-noxious proteins can be produced without deleterious effects on the plant.

10 One significant economic area of interest is the use of transgenic plants for pest control.

Pests such as insects, nematodes and mites are a significant economic cost to plant-based industries. Losses arise through production lost to pest consumption, spoilage and introduction of disease carried by pests.

15 Traditionally, control of pests has been pursued through the application of pesticidal chemicals. Continued use of chemicals is subject to a number of disadvantages. Pests can develop tolerance to chemicals over time producing pesticide resistant populations. Chemical residues may also pose environmental hazards as well as health concerns.

20 Biological control presents an alternative means of pest control which is potentially more effective and specific than current methods, as well as reducing dependence on chemical pesticides. The need for biological controls has lead to the use of recombinant DNA techniques to insert genes which express pesticidal toxins into plant cells.

25 This technology in turn may also give rise to resistant pest populations. There is therefore an ongoing need to find proteins with pesticidal properties, particularly those that are encoded by single genes. These genes can be used to transform plants to produce pest resistant cultivars.

30 Genes studied to date include a range of *cry* genes from the bacterium *Bacillus thuringiensis* (Bt) encoding  $\beta$ -endotoxins and various higher plant genes encoding antimetabolites such as protease and  $\alpha$ -amylase inhibitors and lectins (Boulter, 1993). Many transgenic cultivars with improved insect resistance are now being commercialised, for example, 35 transgenic cotton, corn, and potatoes (James and Krattiger, 1996).

The commercial production of avidin from reproductive tissue of plants using such constructs has also been contemplated (US 5,767,379). The production methods are subject to a number of drawbacks. Male fertility in plants can be lost and expression in vegetative 40 tissue may be low. This may be due in part to expression being outside the cell.

Most recently, the use of avidin and streptavidin as larvicides against insect pests has been explored (WO 94/00992; Morgan *et al.*, 1993; and Bruins *et al.*, *Insect Biochemistry*, 21: 535-539, 1991). In WO 94/00992 generation of resistant plants has been sought by inserting into the cells of a plant a gene whose expression causes production of one or  
5 more of those glycoproteins in larvicidal amounts. While transient expression was shown in maize cells in suspension, no data is presented to show that avidin or streptavidin were expressed at insecticidal concentrations or that plants could be produced expressing same without deleterious side effects.

10 In later applications by the same applicant as for WO 94/00992, transgenic plants with avidin under control of a promoter are described, see WO 96/40949, WO 99/04023 and US 5,767,379. There is no mention of any of the plants produced in these documents as having insecticidal activity. Moreover, the plants produced all exhibit male sterility. There is no specific suggestion in these documents that vacuole targeting could be used  
15 to avoid development of male sterility. Similarly, in *Plant Physiol.* 102 (Suppl.): 45, 1993 a chimeric gene comprising streptavidin coding sequences under control of the CaMV 35S promoter and three signal sequences is contemplated. The signal sequences are indicated as useful for targeting protein to different organelles in plants. However, these organelles are unspecified. Moreover, there is no evidence any plants have been produced incorporating  
20 the chimeric genes nor any discussion as to the effects the genes may have on those plants.

The issues with chimeric genes is whether they can be correctly targeted, whether they will be stable in vacuoles, and whether sequestration in a cell vacuole will prevent the  
25 protein expressed by the chimeric gene from having deleterious effects on the plant cells.

To date, limited success has been achieved in producing insect resistant plants using this technology.

30 Specifically, no one has been able to produce a fertile plant expressing significant levels of a biotin-binding protein in vegetative tissues, nor plants shown to be resistant to insect attack due to the expression of a biotin-binding protein. Similarly, no one has yet been able to provide a protein conferring broad spectrum insect resistance on a host plant without deleterious effects to the plant.

35 It is an object of the present invention to provide chimeric polypeptides and plants which go some way to overcoming the above drawbacks or at least to provide the public with a useful choice.

40

## SUMMARY OF THE INVENTION

Accordingly, in one aspect, the present invention may be broadly said to consist in a chimeric polypeptide that comprises (a) a vacuole targeting sequence encoding a polypeptide; and  
5 (b) a sequence encoding a plant-noxious protein linked in operable combination to said targeting polypeptide.

Preferably, the vacuole targeting polypeptide is a signal sequence polypeptide selected from proteinase inhibitor signal sequence (PPI-I or PPI-II) polypeptide which have the  
10 amino acid sequences set out in Figure 8B and Figure 9B respectively, or variants thereof having substantially equivalent signalling activity thereto.

Preferably, the plant-noxious protein is pest control protein and desirably, a biotin-binding protein.  
15

Preferably, the biotin-binding protein encoded is avidin or streptavidin or a functionally equivalent variant thereof.

The chimeric polypeptides may further comprise at least one additional sequence encoding a protein or peptide.  
20

Conveniently, the chimeric polypeptides of the invention are obtained by expression of a DNA sequence encoding the chimeric polypeptide in a host cell or organism.

25 In a further aspect, the present invention provides an isolated nucleic acid molecule encoding a chimeric polypeptide of the invention.

This nucleic acid molecule can be an RNA or cDNA molecule but is preferably a DNA molecule.  
30

Also provided by the present invention are recombinant expression vectors which contain a DNA molecule of the invention, and hosts transformed with the vector of the invention capable of expressing a polypeptide of the invention.

35 In a still further aspect, the invention provides a method of producing a polypeptide of the invention comprising the steps of:

- (a) culturing a host cell which has been transformed or transfected with a vector as defined above to express the encoded polypeptide of the invention; and optionally  
40
- (b) recovering the expressed polypeptide.

An additional aspect of the present invention provides a ligand that binds to a polypeptide of the invention. Most usually, the ligand is an antibody or antibody binding fragment.

5 In a further aspect, the present invention provides a method for producing a pest resistant plant, comprising transforming the plant genome to include at least one DNA molecule of the invention which includes a sequence encoding a pest control protein.

10 Also provided is a transgenic plant expressing insecticidally effective concentrations of a pest control protein.

The present invention further provides a transgenic plant that contains a DNA molecule of the invention.

15 In one embodiment the transgenic plant further contains at least one additional DNA sequence encoding a protein or peptide.

20 In a still further aspect, the present invention provides a method for controlling or killing pests comprising administering to said pest an amount of a chimeric polypeptide of the invention, which includes a sequence encoding a pest control protein, effective to control or kill said pest.

25 In one embodiment of the method, the chimeric polypeptide is administered with a second pest control protein, where the combination provides more effective control than administration of the second pest control protein alone.

Usually, the pests are the immature stages of insects, including larvae, grubs, nymphs and instars.

30 In yet a further aspect, the present invention provides a composition comprising a chimeric polypeptide of the invention and a carrier, diluent, excipient or adjuvant.

In a further composition aspect, the present invention provides a composition comprising plant material produced in accordance with the invention and a carrier, diluent, excipient or adjuvant.

35 The composition is preferably a pesticidal composition.

40 In a further aspect, the present invention provides a method for controlling or killing pests comprising administering to said pest plant material produced in accordance with the invention, which expresses a pest control protein, or administering a pesticidal composition of the invention, effective to control or kill said pest.

In a still further aspect, the present invention provides a method for producing a plant-noxious protein, the method comprising extracting the protein from a plant containing a DNA molecule of the invention coding for same.

- 5 While the invention is broadly as defined above, it will be appreciated by those persons skilled in the art that it is not limited thereto and that it also includes embodiments of which the following description gives examples.

Figure 1 shows the nucleic acid sequence of Potato Proteinase Inhibitor I (PPI-I/pUC19). The signal sequence is in bold type and the start and stop codons are in italic. The mutagenic primer is denoted by underlined in lower case with the Bgl II site created by mutagenesis in bold italic. The upstream and downstream primers used were the Forward and Reverse M13(lacZ) Primers [Perkin Elmer].

15 Figure 2 shows Avidin cDNA (pGEMav). The signal sequence represented in bold type, start and stop codons are in italic, primers are underlined lower case with the BamH I site created by mutagenesis in italic. The downstream primer used was the Reverse M13(lacZ) Primer [Perkin Elmer].

20 Figure 3 shows streptavidin cDNA (Streptavidin/pUC19). Start and stop codons are in bold type. EcoR I and Xba I sites are in italic.

25 Figure 4 shows potato proteinase inhibitor II (PPI-II/pUC19). The signal sequence is represented in bold type and start and stop codons are in bold italic. Underlined type denotes the intron within the signal sequence. The asterisk denotes the result of PCR error during isolation of the PPI-II sequence.

30 Figure 5 shows components of the ligation reaction to produce recombinant pART7 containing the PPI-I signal sequence/Avidin cDNA gene fusion. A) PPI-I leader fragment resulting from a Sal I/Bgl II digest of the mutated PPI-I PCR product. B) Avidin mature protein cDNA fragment, resulting from a BamH I/Hind III digest of the mutated Avidin PCR product. C) pART7 vector following an Xho I/Hind III digestion. \* denotes compatible cohesive ends. \*\* denotes compatible cohesive ends.

35 Figure 6 shows DNA fragments A, B and C were the components of the ligation reaction to produce recombinant pUC19 containing the PPI-II signal sequence/Streptavidin cDNA gene fusion. The fused gene was then released from pUC19 by a Sal I/BamH I digest and ligation of components D and E produced recombinant pART7. A) PPI-II leader fragment resulting from a Sal I/EcoR I digest of the PPI-II PCR product. B) Streptavidin cDNA  
40 fragment, resulting from an EcoR I/Xba I digest of the recombinant plasmid pUC19/Streptavidin cDNA. D) PPI-II signal sequence/Streptavidin cDNA gene fusion

fragment, resulting from a Sal/BamH I digest of recombinant pUC19 containing the fused gene. E) pART7 vector following an Xho I/BamH digestion. \* denotes compatible cohesive ends.

5 Figure 7 shows a schematic representation of the pART7 expression cassette as it was cloned into the pART27 binary vector; A) containing the PPI-I-Avidin gene fusion and B) containing the PPI-II/Streptavidin gene fusion.

10 Figure 8 shows PPI-I/Avidin gene fusion sequence (A) and fusion protein sequence (B): The fusion protein has a total of 161 amino acids; the PPI-I sequence is represented by italic type with bold type denoting the PPI-I signal peptide. Two amino acids, novel to both the PPI-I and the Avidin peptide sequences and represented in lower case were introduced with the ligation of the Bgl II and BamH I compatible cohesive ends.

15 Figure 9 shows PPI-II/Streptavidin gene fusion sequence (A) and fusion protein sequence (B): The fusion protein has a total of 168 amino acids; the PPI-II sequence is represented by italic type with bold type denoting the PPI-II signal peptide. Three amino acids, novel to both PPI-II and the Streptavidin peptide sequences and represented in lower case were introduced at the point of fusion.

20 Figures 10 and 11 show the survival of larvae of the potato moth, *Phthorimaea operculella* fed tobacco plants expressing avidin in two replicate trials.

25 Figure 12 (A) shows nucleotide sequence for the gene for streptavidin (Argarana *et al.* 1986). The signal sequence is represented in bold type, start and stop codons in bold italic. (B) shows protein sequence for streptavidin. The signal sequence is represented in bold type.

30 Figure 13 shows a cross section of a transgenic leaf stained with methylene blue/Azure II to show general structure of the leaf. Densely stained bodies in the vacuole are arrowed. Bar = 50  $\mu$ m. v; vascular bundle. t; trichome. g; glandular hair.

35 Figure 14 shows immunolabelling of the section for the distribution of avidin (arrowed). Fluorescence indicates the presence of avidin. Bar = 50  $\mu$ m.

Figure 15 shows a transmission electron micrograph showing the distribution of protein bodies in the vacuole of the cell (arrowed). Bar = 1  $\mu$ m.

40 Figure 16 shows a higher magnification of Figure 15. Immunogold labelling over the surface of the protein bodies within the vacuole (arrowed). Bar = 200nm.

Figure 17 shows the survival of larvae of the potato tuber moth, *Phthorimaea operculella* fed tobacco plants expressing streptavidin in two replicate trials.

5 Figure 18 shows the proportion of larvae of the potato tuber moth, *Phthorimaea operculella* at each instar after feeding for nine days tobacco plants expressing streptavidin.

Figure 19A shows the growth of larvae of the common cutworm, *Spodoptera litura*, fed tobacco leaves expressing avidin.

10 Figure 19B *Spodoptera litura* larvae used in Example 8 are shown on Day 15 of the trial. Larvae fed control tobacco are pictured on the left, and larvae fed tobacco expressing avidin are on the right.

15 Figure 19C *Spodoptera litura* larvae used in Example 8 are shown on Day 15 of the trial, in boxes with tobacco leaves used in the experiment. A typical control treatment with large larvae and stripped leaves is shown on the left, a typical avidin-fed treatment with small dead larvae and minimally damaged leaves on the right.

20 Figure 20 shows the survival of larvae of the common cutworm, *Spodoptera litura*, fed tobacco leaves expressing avidin.

Figure 21 shows the accumulation of larval biomass of the common cutworm, *Spodoptera litura*, fed tobacco leaves expressing avidin.

25 Figure 22A shows the growth of larvae of the cotton bollworm (corn earworm, tomato fruitworm), *Helicoverpa armigera* fed tobacco leaves expressing avidin.

30 Figure 22B *Helicoverpa armigera* larvae used in Example 8 are shown on Day 14 of the trial. Larvae fed control tobacco are pictured on the left, and larvae fed tobacco expressing avidin are on the right.

35 Figure 22C *Helicoverpa armigera* larvae used in Example 8 are shown on Day 14 of the trial, in boxes with tobacco leaves used in the experiment. A typical control treatment with large larvae and stripped leaves is shown on the left, a typical avidin-fed treatment with small dead larvae and minimally damaged leaves on the right.

Figure 23 shows the survival of larvae of the cotton bollworm (corn earworm, tomato fruitworm), *Helicoverpa armigera*, fed tobacco leaves expressing avidin.

40 Figure 24 shows the accumulation of biomass of larvae of the cotton bollworm (corn earworm, tomato fruitworm), *Helicoverpa armigera*, fed tobacco leaves expressing avidin.



Figure 25 shows the effect of the level of avidin expression in tobacco on the growth of larvae of the cotton bollworm (corn earworm, tomato fruitworm), *Helicoverpa armigera*.

- 5 Figure 26 shows the effect of the level of avidin expression in tobacco on the survival of larvae of the cotton bollworm (corn earworm, tomato fruitworm), *Helicoverpa armigera*.

Figure 27 shows the effect of the level of avidin expression in tobacco on the accumulation of biomass of larvae of the cotton bollworm (corn earworm, tomato fruitworm), *Helicoverpa armigera*.

10

Figure 28 shows the effect of avidin and streptavidin incorporated into insect diet at three concentrations on the growth of larvae of the pine shoot tip moth, *Rhyacionia buoliana*.

- 15 Figure 29 shows the effect of avidin and streptavidin incorporated into insect diet at three concentrations on the survival of larvae of the pine shoot tip moth, *Rhyacionia buoliana*.

Figure 30 shows the effect of avidin and streptavidin incorporated into insect diet at three concentrations on the accumulation of biomass of larvae of the pine shoot tip moth, *Rhyacionia buoliana*.

20

Figure 31 shows the effect of avidin-painted willow leaves on the survival of larvae of the willow sawfly, *Nematus oligospilus*.

- 25 Figure 32 shows the effect of avidin-painted willow leaves on the weight gain of larvae of the willow sawfly, *Nematus oligospilus*.

Figure 33 shows the effect of avidin-painted willow leaves on the formation of the pupae of the willow sawfly, *Nematus oligospilus*.

- 30 Figure 34 shows the effect of avidin-painted willow leaves on the emergence of adults of the willow sawfly, *Teleogryllus commodus*.

Figure 35 shows the effect of avidin-painted lettuce leaves on the growth of nymphs of the black field cricket, *Teleogryllus commodus*.

35

Figure 36 shows the effect of avidin-painted lettuce leaves on the survival of nymphs of the black field cricket, *Teleogryllus commodus*.

- 40 Figure 37 shows the effect of avidin-painted lettuce leaves on the accumulation of biomass of nymphs of the black field cricket, *Teleogryllus commodus*.

Figure 38 shows the effect of streptavidin incorporated into insect diet on the survival of neonate larvae of the clover root weevil, *Sitona lepidus*.

- 5 Figure 39 shows the effect of streptavidin incorporated into insect diet on the survival of larvae of the Argentine stem weevil, *Listronotus bonariensis*.

Figure 40 shows the effect of avidin-painted clover leaves on the survival of adults of the clover root weevil, *Sitona lepidus*.

10

Figure 41 shows the effect of avidin added to pollen on the consumption of that food by adult honeybees, *Apis mellifera*.

15

Figure 42 shows the effect of avidin added to pollen on the survival of adult honeybees, *Apis mellifera*.

Figure 43 shows the effect of avidin-painted lettuce leaves on the weights of snails, *Cantareus aspersus*.

20

Figure 44 shows the effect of avidin-painted lettuce leaves on the survival of snails, *Cantareus aspersus*.

Figure 45 shows the effect of avidin-painted lettuce leaves on the weight of slugs, *Deroceras reticulatum*.

25

Figure 46 shows the effect of avidin-painted lettuce leaves on the survival of slugs, *Deroceras reticulatum*.

30

Figure 47 shows the effect of avidin expression in tobacco combined with painted-on aprotinin or Cry1Ba on survival of larvae of the cotton bollworm (corn earworm, tomato fruitworm), *Helicoverpa armigera*.

35

Figure 48 shows the effect of avidin expression in tobacco combined with painted-on aprotinin on growth of larvae of the cotton bollworm (corn earworm, tomato fruitworm), *Helicoverpa armigera*.

40

Figure 49 shows the effect of avidin expression in tobacco combined with painted-on aprotinin on biomass of larvae of the cotton bollworm (corn earworm, tomato fruitworm), *Helicoverpa armigera*.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel chimeric polypeptides comprising vacuole targeting sequences and plant-noxious sequences. The targeting sequences and plant-noxious sequences are operably linked.

The term "operably linked" as used herein refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. For example, a signal sequence is operably linked to a coding sequence if the promoter affects its transcription or expression.

The term "vacuole targeting sequence" as used herein refers to a sequence operable to direct or sort a selected non-vacuolar protein to which such sequence is linked, to a plant vacuole.

The vacuolar targeting polypeptide sequences of the invention, when transformed into plants, function to direct or sort the protein products directed by the expression of genes to which they are operably linked from the cytoplasm to the vacuole of the plant cell. Since the vacuole of plant cells has a storage function, proteins directed there remain there, continually increasing in abundance, unless subject to degradation by vacuolar proteinases. The vacuolar proteins are also isolated from the major metabolic processes in the plant and thus will not interfere with the plant growth and development. The success of the present invention needed that both these requirements be met.

Vacuolar targeting sequences include any such targeting sequences as are known in the art that effect proper vacuole targeting in plant hosts. These include polypeptides targeting barley lectin (Bednarek *et al.*, 1990), sweet potato sporamin (Matsuoka *et al.*, 1990), tobacco chitinase (Neuhaus *et al.*, 1991), bean phytohemagglutinin (Tague *et al.*, 1990), 2S albumin (Saalbach *et al.*, 1996), aleurain (Holwerda *et al.*, 1992). Vacuolar targeting in plants has been widely studied (for example see Chrispeels, 1991; Chrispeels & Raikhel, 1992; Dromboski & Raikhel, 1996; Kirsch *et al.*, 1994; Nakamura & Matsuoka, 1993; Neilsen *et al.*, 1996; Rusch & Kendall, 1995; Schroder *et al.*, 1993; Vitale & Chrispeels, 1992; von Heijne, 1983). Other sequences are described, for example, in US 5,436,394, US 5,792,923, US 5,360,726, US 5,525,713 and US 5,576,428 incorporated herein by reference. However, potato proteinase inhibitor targeting sequences are preferred.

A number of potato proteinase signal sequence polypeptides designated PPI-I and PPI-II are disclosed for use herein. These polypeptides were described previously (Beuning *et al.*, (1994); Christeller *et al.* (1994)). The polypeptides have the amino acid sequences set out in Figures 8B and 9B respectively. Also encompassed within the invention are variants of these polypeptides and those known in the art which have substantially equivalent

targeting sequence activity thereto.

5 The term "variant" as used herein refers to a polypeptide wherein the amino acid sequence exhibits substantially 70% or greater homology with the amino acid sequences set out in Figures 1 and 4. Preferably, the variants will have greater than 85% homology, and most preferably, 95% homology or more. Variants may be arrived at by modification of the native amino acid sequence by such modifications as insertion, substitution or deletion of one or more amino acids.

10 As noted above, the chimeric polypeptide comprises a vacuole targeting signal sequence operably linked to a plant-noxious protein.

15 The term "plant-noxious protein" as used herein refers to a protein which has a negative effect on plant health, growth, development or fertility when not sequestered in a plant vacuole.

20 Examples of plant-noxious proteins include barnase (ribonuclease), cellulases and other cell wall degrading enzymes such as pectinases and polygalacturonases as well as pest control proteins discussed below.

25 In one embodiment, the plant-noxious protein is a pest control protein. Pest control proteins include proteins which decrease availability of vitamins, or other essential growth component or are toxic to pests *per se*. Toxic proteins include lectins, proteinase inhibitors, *Bacillus thuringiensis* insecticidal proteins, alpha-amylase inhibitors, vegetative insecticidal proteins, lipoxxygenase and cholesterol oxidase. Proteins which decrease availability of vitamins fall broadly into these categories of degradative enzymes and binding proteins. Examples of degradative enzymes include thiaminase, riboflavin hydrolase, and pantothenate hydrolase but are not limited thereto.

30 Bt proteins useful in the present invention include Cry proteins such as Cry1Ba, Cry1Ac, Cry1Cb, Cry1Da, Cry1F, Cry5 and Cry9A, but are not limited thereto.

35 Proteinase inhibitors useful in the invention include aprotinin, kunitz-type inhibitors from soybean, arrowroot, taro, proteinase inhibitor 1, proteinase inhibitor 2, alpha-1 antitrypsin, Bownan-Birk inhibitors from soybean and cowpea and oryzacystatin.

40 The term "pest" as used herein refers to a broad group of organisms which at some point in their life cycle live or feed on plants adversely affecting same. Included in the term are protozoa, arthropods (especially insects), aschelminthes and platyhelminthes, nematodes and molluscs.

Binding proteins useful in the invention include riboflavin-binding protein, carotenoid binding proteins, fatty-acid binding proteins, retinol binding proteins, alpha-tocopherol binding proteins, folate-binding proteins, thiamine-binding proteins, pantothenate-binding proteins and biotin-binding proteins, but again are not limited thereto. A preferred group of binding proteins are vitamin binding proteins, particularly biotin-binding proteins. These are proteins which associate with biotin to form a complex with a dissociation constant of  $10^{-6}$ M or less. Usually, the complex is a non-covalent complex. The biotin binding proteins for use herein must be operable to bind biotin in a plant system without adversely affecting the plant, or to affect the plant in a minimal way, when included in chimeric polypeptides of the invention. For example, slight reductions in plant growth would be acceptable.

Systems requiring covalent enzymatic sequestration are also contemplated within this term. For example, simultaneous overexpression of a biotin requiring carboxylase or a biotin acceptor peptide (for example, see Schatz, P.J., *Biotechnology*, 11: 1138-1143, 1993) and biotin holocarboxylase synthetase in the vacuole could be used to induce biotin deficiency. Biotin would be covalently sequestered enzymatically on vacuole rupture.

Biotin is an essential nutrient for many species of pests (Dadd, R.H., 1985; Kerkut G.A. *et al.*, *Comprehensive Insect Physiology, Biochemistry and Pharmacology*, 4: 313-390, 1985). As discussed above, biotin-binding proteins have been found to have pesticidal properties and to inhibit growth of pests. The binding of biotin causes a biotin deficiency which results in the inhibition of growth and ultimate death of pests.

Biotin-binding proteins known in the art include egg yolk biotin-binding proteins (Subramanian and Ariga, 1995, *Biochem. J.*, 308: 573-577, serum (Seshagiri and Ariga, 1987, *Biochem. Biophys. Acta*, 916: 474-481), biotin-binding antibodies, and fragments thereof, biotin holocarboxylase synthetase, biotinidase, bacterial proteins, avidin, isolated from egg white, and streptavidin. The properties of a number of these proteins are usefully discussed in *Methods of Enzymology* Vol 184 (eds M. Wilcheta and E A Bayer).

Preferred biotin-binding polypeptides, for use in the present invention, are avidin and streptavidin or functionally equivalent variants thereof. It will be appreciated that other groups that function to bind biotin, such as those referred to above, are equally able to be used in the present invention.

Avidin is a water-soluble tetrameric glycoprotein isolated originally from raw egg white (*J. Biol. Chem.* 136: 801 (1940)). The protein is well known with the complete amino acid sequence having been published in, for example, *J. Biol. Chem.* 246: 698 (1971). The full amino acid sequence for avidin is shown in Figure 8B (amino acids 34 to 161). Several natural variants of avidin have also been discussed in Keinänen *et al.*, *Eur. J. Biochem.*

220:615-621 (1994) and synthetic variants in Marttila *et al.*, FEBS Letters. 441:313-317 (1998).

Streptavidin is a non-glycosylated bacterial binding protein derived from the culture supernatant of *Streptomyces avidinii* (Bayer *et al.*, 1990). The full amino acid sequence for streptavidin is given in Figure 12.

'Core' SAV is equivalent to amino acid residues 37-164 of *Streptomyces avidinii* (SAV) Figure 12, (Argarana *et al.*, 1986). Other 'core' SAV molecules have been produced with various N-terminal and C-terminal deletions. A preferred sequence referred to as "Synthetic 'Core' Streptavidin" is a modified 'core' SAV having the sequence shown in Figure 9B (amino acids 41 to 168). SYNSAV is equivalent to 'Core' SAV modified such that codons for each amino acid correspond to those in highly expressed *E.coli* genes. SYNSAV is also modified to contain unique restriction sites evenly throughout sequence. The resulting sequence has G + C content of 54% relative to 69% for same region of native SAV (Thompson *et al.* (1993))<sup>29</sup>. A number of natural variants of streptavidin have also been described in Bayer *et al.*, *Biochem. Biophys. Acta* 1263: 60-66 (1995), GenBank Acc. No. S78782 and S78777. Synthetic streptavidin molecules can also be produced using known art techniques. See for example WO 89/03422.

The chimeric polypeptides of the invention may further comprise one or more sequences encoding other proteins or peptides. Two to four further sequences are contemplated, but more are feasible. These other proteins or peptides may be selected from any proteins known in the art which it is desired to express in a plant vacuole including plant-noxious proteins discussed above.

Proteins to be produced in conjunction with pest control proteins may be selected so as to achieve an additive or synergistic effect as demonstrated in Example 18), a broader spectrum of control, or to reduce the risk of resistance developing. Examples of such proteins include other pest control proteins as discussed above including proteinase inhibitors, toxic proteins and biotin-binding proteins, as well as antimicrobial, antifungal and antiviral proteins but not limited thereto.

The applicants have surprisingly found that plants expressing avidin when combined with Bt insecticidal protein can exhibit synergistic effects on pests (Figure 47). Proteinase inhibitors may be desirable for use in preventing proteolysis of the insect control protein (see Example 18) Shao *et al.*, *J Invertebr. Pathol.* 72: 73-81 (1998); and Keller *et al.*, *Insect Biochem. Mol. Biol.* 26: 365-73 (1996). The compatibility of biotin-binding proteins and protease inhibitors has been demonstrated by the applicant.

The antimicrobial, antifungal and antiviral groups of proteins can assist in the control

of plant disease particularly where insect damage contributes to the spread of disease. Proteins which have been shown to have these activities include dermaseptins, cercropins, attacins, lysozyme, chitinases, hevein, glucose oxidase, glucanases, thionins, lectins, *Raphanus sativus*, antifungal protein, osmotin, lipid transfer proteins, lipoxygenase and virus coat proteins.

Similarly, reduction in disease from insect resistant crops has been reported. For example, research at Iowa State University has shown reduction in feeding damage is linked to a reduction in ear mould in Bt maize.

The reader will appreciate that modifications, including chemical and biochemical modifications, of the polypeptides of the invention are possible. Such modifications include, for example, acetylation, carboxylation, phosphorylation, glycosylation, ubiquitination, labelling, and the like. The production of peptide fragments is also well within the capabilities of an art skilled worker.

The polypeptides of the invention can be prepared in a variety of ways. For example, as indicated above the signal sequences and biotin-binding proteins can be produced by isolation from natural sources and then coupled using techniques known in the art. For example, through recombinant nucleic acid methods.

Synthesis using known techniques (such as stepwise solid phase synthesis described by Merryfield, *J. Amer. Chem. Soc.* Vol 85:2149-2156, 1963), or as preferred through employing recombinant DNA techniques.

Variants of the polypeptide can similarly be made by any of those techniques known in the art. For example, variants can be prepared by site-specific mutagenesis of the DNA encoding the native amino acid sequence as described by Adelman et al. *DNA* 2:183 (1983). Generally, the variants produced are functionally equivalent to the original sequence.

Where it is preferred, recombinant techniques used to produce the polypeptide of the invention, the first step is to obtain DNA encoding the desired product. Such DNA comprises a still further aspect of this invention.

The DNA of the invention may encode a native or modified polypeptide of the invention or an active fragment thereof. In its presently preferred forms, the DNA comprises the nucleotide sequence of Figure 8A, or the nucleotide sequence of Figure 9A. Preferred sequences exhibit 60% or greater homology with these sequences, preferably 80% homology and most preferably 95% homology or more. That is, most preferred sequences will hybridise to the sequences of the invention under stringent hybridisation conditions.

The DNA can be isolated from any appropriate natural source or can be produced as intron free cDNA using conventional techniques. DNA can also be produced in the form of synthetic oligonucleotides where the size of the active fragment to be produced permits. By way of example, the Triester method of Matteucci *et al. J. Am. Chem. Soc.* Vol 103:3185-3191 (1981) may be employed.

Where desirable, the DNA of the invention can also code for a chimeric polypeptide of the invention (including polypeptides encoding more than one protein). Such fusion proteins may be produced as disclosed in WO 86/02077 incorporated herein by reference. Fusion proteins further comprising the polypeptide of the invention and a carrier protein are possible. This carrier protein will generally be cleavable from the polypeptide, peptide or fragment under controlled conditions. Examples of commonly employed carrier proteins are  $\beta$ -galactosidase and glutathione-S-transferase.

As indicated above, also possible are variants of the polypeptide or peptide which differ from the native amino acid sequence by insertion, substitution or deletion of one or more amino acids. Neutral variations (those which have no effect on function) are specifically contemplated. Where such a variant is desired, the nucleotide sequence of the native DNA is altered appropriately. This alteration can be made through elective synthesis of the DNA or by modification of the native DNA by, for example, site-specific or cassette mutagenesis. Preferably, where portions of cDNA or genomic DNA require sequence modifications, site-specific primer directed mutagenesis is employed using techniques standard in the art.

In a further aspect, the present invention consists in replicable transfer vectors suitable for use in preparing a polypeptide or peptide of the invention. These vectors may be constructed according to techniques well known in the art, or may be selected from cloning vectors available in the art.

The cloning vector may be selected according to the host or host cell to be used. Useful vectors will generally have the following characteristics:

- (a) the ability to self-replicate;
- (b) the possession of a single target for any particular restriction endonuclease; and
- (c) desirably, carry genes for a readily selectable marker such as antibiotic resistance or herbicide tolerance.

Two major types of vector possessing these characteristics are plasmids and bacterial viruses (bacteriophages or phages). Presently preferred vectors include the plasmids pMOS-Blue, pGem-T, pUC18, pUC19, pART27, pMON, pJIT, pBIN, pRD 400, pART7.

Also contemplated is the use of RNA vectors for example, tobacco mosaic virus (Donson *et al., Proc Natl. Acad. Sci. USA.*, 88:7204-8, 1991), potato virus X (PVX) (Chapman



*et al.*, *Plant J.* 2:549-57, 1992), and barley stripe mosaic virus (ESMV) (Josh, *et al.*, *EMBO J.* 9:2663-9, 1990). TMV has previously been used to infect plants to produce therapeutic protein products (Turpen, *Philos Trans. R. Soc. Lond. Biol. Sci.*, 354: 665-73, 1999). Basic RNA vectors can be produced according to known art techniques.

5 The DNA molecules of the invention may be expressed by placing them in operable linkage with suitable control sequences in a replicable expression vector. Control sequences may include origins of replication, a promoter, enhancer and transcriptional terminator sequences amongst others. The selection of the control sequence to be included in the expression  
10 vector is dependent on the type of host or host cell intended to be used for expressing the DNA.

15 Generally, procaryotic, yeast, insect or mammalian cells are useful hosts. Also included within the term hosts are plasmid vectors. Suitable procaryotic hosts include *E. coli*, *Bacillus* species and various species of *Pseudomonas*. Commonly used promoters such as  $\beta$ -lactamase (penicillinase) and lactose (*lac*) promoter systems are all well known in the art. Any available promoter system compatible with the host of choice can be used. Vectors used in yeast are also available and well known. A suitable example is the 2 micron origin of replication plasmid.

20 Similarly, vectors for use in mammalian cells are also well known. Such vectors include well known derivatives of SV-40, adenovirus, retrovirus-derived DNA sequences, *Herpes simplex* viruses, and vectors derived from a combination of plasmid and phage DNA.

25 Further eucaryotic expression vectors are known in the art (e.g. P.J. Southern and P.Berg, *J. Mol. Appl. Genet.* 1 327-341 (1982); S. Subramani *et al.*, *Mol. Cell. Biol.* 1, 854-864 (1981); R.J. Kaufmann and P.A. Sharp, "Amplification and Expression of Sequences Cotransfected with a Modular Dihydrofolate Reducase Complementary DNA Gene, *J. Mol. Biol.* 159, 601-621 (1982); R.J. Kaufmann and P.A. Sharp, *Mol. Cell. Biol.* 159, 601-  
30 664 (1982); S.I. Scahill *et al.*, "Expressions And Characterization Of The Product Of A Human Immune Interferon DNA Gene In Chinese Hamster Ovary Cells," *Proc. Natl. Acad. Sci. USA.* 80, 4654-4659 (1983); G. Urlaub and L.A. Chasin, *Proc. Natl. Acad. Sci. USA.* 77, 4216-4220, (1980).

35 The expression vectors useful in the present invention contain at least one expression control sequence that is operatively linked to the DNA sequence or fragment to be expressed. The control sequence is inserted in the vector in order to control and to regulate the expression of the cloned DNA sequence. Examples of useful expression control sequences are the *lac* system, the *trp* system, the *tac* system, the *trc* system, major operator and promoter  
40 regions of phage lambda, the glycolytic promoters of yeast acid phosphatase, e.g. *Pho5*, the promoters of the yeast alpha-mating factors, and promoters derived from polyoma,

adenovirus, retrovirus, and simian virus, e.g. the early and late promoters of SV40, and other sequences known to control the expression of genes of prokaryotic and eucaryotic cells and their viruses or combinations thereof.

5 Also useful in the present invention are promoters which can be used to target proteins to specific plant tissues. These have application in situations where accumulation of a protein in a particular tissue is desired, or alternatively, is to be avoided to prevent non-target effects. For example, accumulation of an insect control protein in pollen may be undesirable if it is fed on by non target pests such as butterflies, bees or other pollinators. Specific  
10 promoters can be used to target such pest control proteins away from pollen.

Alternatively, a target pest may have defined feeding characteristics such as only feeding on leaves, seed, fruit, flowers or the like. In such cases, it would be desirable to target the pest control protein to the plant tissues being feed on, or to particular cells within  
15 those tissues. For example, to leaf epidermal cells, root cortex cells, mesophyll cells and the like. Any of the promoters known in the art for targeting specific plant tissues may be employed.

Preferred promoters for use herein include lacZ, CaMV-35S, LHC a/b, T7, nos, rubisco small subunit (SSU), gpd and nod gene promoters.  
20

In the construction of a vector it is also an advantage to be able to distinguish the vector incorporating the foreign DNA from unmodified vectors by a convenient and rapid assay. Such assays include measurable colour changes, antibiotic resistance, herbicide tolerance  
25 and the like. In one preferred vector, the  $\beta$ -galactosidase gene is used, which gene is detectable by clones exhibiting a blue phenotype on X-gal plates. This facilitates selection.

Once selected, the vectors may be isolated from the culture using routine procedures such as freeze-thaw extraction followed by purification.  
30

For expression, vectors containing the DNA of the invention to be expressed and control signals are inserted or transformed into a host or host cell. Intermediate host cells can be used to increase the copy number of the cloning vector prior to introduction into plant cells. Some useful expression host cells include well-known prokaryotic and eucaryotic  
35 cells. Some suitable prokaryotic hosts include, for example, *E. coli*, such as *E. coli*, S G-936, *E. coli* HB 101, *E. coli* W3110, *E. coli* X1776, *E. coli*, 2282, *E. coli*, DHT, and *E. coli*, MR01, *Pseudomonas*, *Bacillus*, such as *Bacillus subtilis*, and *Streptomyces*. Suitable eucaryotic cells include yeast and other fungi, insect, animal cells, such as COS cells and CHO cells, human cells and plant cells in tissue culture.

40 Expression systems employing insect cells utilising the control systems provided by baculovirus

vectors have been described (Miller *et al.*, in *Genetic Engineering*, 8: 277-297, 1986).

Depending on the host used, transformation is performed according to standard techniques appropriate to such cells. For prokaryotes or other cells that contain substantial cell walls, the calcium treatment process (Cohen, S. N. *Proceedings, National Academy of Science, USA* 69: 2110, 1972) may be employed. For mammalian cells without such cell walls the calcium phosphate precipitation method of Graeme and Van Der Eb, *Virology* 52:546, 1978 is preferred. Transformations into plants may be carried out using *Agrobacterium tumefaciens* (Shaw *et al.*, *Gene* 23:315, 1983) or into yeast according to the method of Van Solingen *et al. J.Bact.* 130: 946, 1977 and Hsiao *et al. Proceedings, National Academy of Science*, 76: 3829, 1979.

In a preferred transformation process, the vectors of the invention are incorporated into *Agrobacterium tumefaciens* which can be used to infect plant cells, particularly dicotyledenous plant cells, thereby transferring the vectors and conferring pest resistance. The cloning vectors can also be introduced into plant cells using convenient art techniques such as electroporation, microparticle bombardment and microinjection. Microparticle bombardment is the preferred transformation process for monocotyledenous plants. Suitable plant transformation techniques are usefully summarised in Torres *et al.*, *Plant Cell, Tissue and Organ Culture* 34: 279-285, 1993, Michelmore *et al.*, *Plant Cell Reports* 6:439-442, 1987, Horsch *et al.*, *Plant Molecular Biology Manual AS*: 1-9, 1988, Xinrun *et al.*, *J. Genet. and Breed.* 46: 287-290, 1992 and WO 97/17455 incorporated herein by reference.

Upon transformation of the selected host with an appropriate vector the polypeptide encoded can be produced, often in the form of fusion protein, by culturing the host cells. The polypeptide of the invention may be detected by rapid assays as indicated above. The polypeptide can then be recovered and purified if desired. Recovery and purification can be achieved using any of those procedures known in the art, for example by absorption onto and elution from an anion exchange resin. This method of producing a polypeptide of the invention constitutes a further aspect of the present invention.

The present invention also provides a method for producing a plant-noxious protein, the method comprising extracting the protein from a plant incorporating a DNA sequence of the invention coding for same. The expression level of the protein may be increased by further incorporating into the DNA sequence of the invention a peptide export signal sequence, or intron sequence. Methods of enhancing expression levels and methods for production of the protein generally may be effected according to the techniques of WO 97/17455 incorporated herein by reference.

The use of the chimeric polypeptides of the present invention represents an advance over this document because the protein is produced in vegetative tissues (leaves, stems, tubers,

roots) as opposed to the reproductive tissues. In the case of avidin and streptavidin this avoids the negative effect of male sterility.

The method of the present invention is also significantly more effective than the disclosed art method with avidin produced as levels up to two times higher than previously reported. See Examples 9, 18 and 17.

The applicant has also been the first to achieve expression of streptavidin using this methodology (see Example 7). The expression levels are approximately twice those previously reported for avidin in US 5,767,379.

It is also noted that the avidin and streptavidin expression levels were achieved with selfed plants. Higher levels of expression are anticipated where plants expressing high levels of avidin or streptavidin are crossed.

The method can be used for producing proteins from a wide range of plants which produce abundant vegetative material e.g. potatoes, cassava, tobacco, grasses, legumes, and trees rather than being restricted to plants which produce large reproductive structures e.g. maize.

In a further aspect, the present invention provides a transgenic plant that contains a DNA molecule of the invention. The plant is produced according to the procedures detailed above, generally comprising transformation with a vector of the invention.

In one embodiment, the transgenic plant contains at least one, and commonly two to four, additional DNA sequences encoding a protein or peptide. More additional sequences are feasible. The proteins or peptides may be any of those proteins or peptides discussed above for the additional protein or peptide within the chimeric polypeptide. Again, sequences encoding Bt Cry proteins are preferred for incorporation into the plant. Incorporation of the additional sequence(s) for the protein(s) or peptide(s) other than as part of the chimeric polypeptide allows for the independent expression of the chimeric polypeptide and additional protein(s) or peptide(s).

Plants suitable for transformation with the vectors of the invention may be selected from a broad range of plants including cereal crops, vegetable, fruit and other food crops, forage crops and turf plants, fibre crops, timber and pulp and paper plants, shelter-belt plants and tree crops, ornamental and flower plants, culinary plants, medicinal plants and herbs and plants grown to produce beverages and plants grown for smoking.

Examples of cereal crops include wheat, rice, barley, maize, oats, millet, sorghum and rye.

Examples of vegetable, fruit and other food crops include root crops such as potato, sweet potato, beetroot, parsnip, turnip, swede and carrot, cucurbits such as cucumbers, pumpkins, squash, marrow, courgettes and watermelon, brassicas such as cauliflower, cabbage, oilseed rape, brussels sprouts and broccoli, corn, tomato, lettuce, celery, onions, garlic, legumes such as lentils, green beans, lima beans, haricot beans, red kidney beans, kudzu beans, mung beans, broadbeans, soybeans, chickpeas, peas, and peanuts, apple, pear, kiwifruit, tamarillo, feijoa apricot, plum, citrus such as orange, lemon, tangelo, grapefruit, uglifruit and mandarin, pineapple, peach, nectarine, cherry, berries, olives and sugarcane.

- 10 Examples of forage crops and turf plants include legumes such as clover, alfalfa, lotus, trefoil and lucerne and grasses and other graminaceous plants such as ryegrass, browntop, fescue, cocksfoot, kikuyu and, paspalum, and sorghum grass.

Fibre crops include cotton, flax, kapok and hemp.

15 Timber, shelterbelt, conservation, pulp and paper plants and tree crops include, for example, pine, eucalyptus, spruce, fir, oak, ash, birch, beech, mahogany, rosewood, ebony, maple, teak, cedar, redwood, jarrah, chestnut, walnut, macadamia nut, poplar, willow, cypress, camphor, mulberry, marram grass and rubberplant.

20 Ornamental shrubs, trees and flower plants include roses, petunias, orchids, carnations, chrysanthemums, daisies, tulips, lilies, gypsophylla, hibiscus, rhododendrons, conifers, camellias, hebes, lavender, lupins, tussock, ferns and native plants.

- 25 Culinary plants include herbs such as basil, rosemary, oregano, bay, and spices such as cinnamon, mace, tumeric, and sage.

Medicinal plants include poroporo, opium poppies, coca, marijuana, camomile, comfrey, foxglove and belladonna.

- 30 Plants used to produce beverages include tea, coffee, hops and cocoa.

Plants used for smoking include tobacco.

- 35 Plants transformed with the vectors of the invention direct expression of the plant-noxious proteins in the vacuoles of the plant cells. The protein is effectively sequestered into the vacuole. Where the protein is a pest control protein, when a pest feeds on the plant expressing a pest control protein, the plant cell components mix together allowing a substance to be controlled (e.g. biotin) to be bound by the binding protein, or alternatively degraded  
40 by enzyme (e.g. in the case of thiamine). This essentially deprives the pest of the vitamin it requires leading to stunted growth and death.

The effect of biotin deprivation is often manifested in the failure of the immature stages of the pests to complete the process of moulting from one developmental stage to the next as demonstrated in Examples 6 to 13.

- 5 In a further aspect, the invention also provides a transgenic plant expressing pesticidally effective concentrations of pest control protein.

- In one preferred aspect, there is provided a plant expressing insecticidally effective concentrations of a biotin-binding protein. Also provided are plants expressing combinations  
10 of biotin-binding proteins and other pest control proteins as discussed above.

The present invention has application in producing plants resistant to a broad range of pests in the larval stage including moths, beetles, weevils, caterpillars, borers, budworms, armyworms, bollworms, rootworms, webworms, aphids, bugs, crickets, locusts, grasshoppers,  
15 grubs, flies, fruitflies, leafminers, plant hoppers, earwigs, scale insects, thrips, and springtails. Plants of the invention may also be resistant to other invertebrate pests of plants such as mites and lice and other pests and pathogens which have a vitamin requirement especially for biotin, particularly those which undergo a moulting process as part of their development.

- 20 List of most preferred pests :

#### Order Lepidoptera:

- cotton bollworm (*Helicoverpa armigera*)
- tropical army-worm (*Spodoptera litura*), also *S. littoralis*, *S. exigua*
- 25 European corn-borer (*Ostrinia nubilalis*)
- tobacco horn worm (*Manduca sexta*)
- loopers (*Chrysodiexis* spp.)
- rice stem borer (*Chilo suppressalis*)
- porina (*Wiseana* spp.)
- 30 cutworms (*Agrotis* spp.)
- diamondback moth (*Plutella xylostella*)
- potato tuber moth (*Phthorimaea operculella*)
- codling moth (*Cydia pomonella*)
- Indian meal moth (*Plodia interpunctella*)
- 35 gypsy moth (*Lymantria dispar*)

#### Order Coleoptera:

- argentine stem weevil (*Listronotus bonariensis*)
- clover root weevil (*Sitona lepidus*)
- 40 grass-grubs (*Costelytra zelandica*, *Odontria* spp.)
- corn rootworm (*Diabrotica virgifera*)

- 23 -

rice and wheat weevils (*Sitophilus* spp.)  
 mealworms (*Tenebrio molitor*)  
 flour beetles (*Tribolium confusum*)

5 Order Orthoptera:

black field cricket (*Teleogryllus commodus*)  
 locusts (*Locusta migratoria*)

Order Hymenoptera:

10 Sawflies (*Sirex* spp., *Nematus olgospilus*)

Order Thysanoptera:

Western Flower thrips (*Frankliniella occidentalis*)

15 Order Diptera:

Hessian flies (*Mayetiola destructor*)

Mites (Class Arachnida)

Order Acari

20 two-spotted mite (*Tetranychus urticae*)  
 European red mite (*Panonychus ulmi*)

The applicants have also demonstrated that plants of the invention will not cause significant mortality of desirable insects such as adult honeybees feeding on pollen (see Example 14). Some specificity of action is also shown where non-moulting, adult stages of insects such as weevils, and invertebrates that do not moult, such as nematodes, slugs or snails, are unlikely to be harmed by feeding on these plants. Hence, plants produced according to the invention have a broad spectrum of pest resistance for invertebrates that moult, particularly insects, as part of their development process.

30 In a further aspect the invention provides a method of imparting pest resistance to plants comprising transforming the plants with a vector according to the present invention.

35 The method may also be effected by transforming isolated plant cells or tissues and generating plants from the transformed cells or tissue using standard culture techniques. Plants at any stage of development, parts thereof, plant cuttings, seeds, plant cells, and cell and tissue cultures transformed with vectors of the invention form further aspects of the invention.

40 Transformed plants can be used in conventional breeding programmes to transfer the DNA sequences of the invention.

The plants of the invention may be grown *en masse*. However, it is also feasible to use a smaller number of plants as "bait" plants within a crop area. Only the bait plants would include the insect control proteins. To ensure preferential targeting of bait plants by pests, attractants such as colour, hormone and scent lures may be used on or around the bait plants.

Alternatively, bait plants may be plants which a target pest has a preference for compared with the crop being grown. For example, it has been shown that rootworm have a preference for *Taiuia* over soybean and maize. Such bait plants may also be used in conjunction with attractants.

In another aspect, the present invention also provides a composition comprising a chimeric polypeptide of the invention and a carrier diluent, excipient or adjuvant therefor. In another composition aspect, there is provided a composition comprising plant material produced in accordance with the invention formulated with agriculturally acceptable excipients, carriers, diluents or adjuvants. The term "plant" as used herein encompasses plants, plant parts such as leaves, roots and flowers, plant cuttings, seeds, tissue cultures, cell cultures and plant cells but is not limited thereto.

Preferably, the composition is a pesticidal composition comprising a pesticidally effective amount of the polypeptide, or plant material and an acceptable carrier. These carriers include inert carriers such as surfactants, spreaders, stickers, mineral and organic granular carriers, stabilisers such as microencapsulation polymers or petroleum-based solvents.

Examples of surfactants, spreaders and stickers include C-Daxoil®, Codacide Oil®, D-C-Trate®, Supawet Oil, Bond®, Boost® Penetrant, Citowett® and Freeway.

Examples of mineral granular spreaders include talcum powder, clay, silica, sand, limestone, gypsum, kaolin, montmorillonite, attapulgite and diatomite.

Examples of organic granular spreaders include corncob granules, pecan shells, peanut hulls and recycled paper fibre.

Examples of stabilisers include sodium tripolyphosphate, UV-absorbers (e.g. 2,4-dihydroxy benzophenone (Uvinul M-400, UM), 4 aminobenzoic acid (PBT), fluorescent brightener-28 (FB-28)), quenchers, radical scavengers, Hindered Amine Light Stabilizers (HALS), photostabilisers (e.g. clays, chromophores) and mineral oils.

Examples of microencapsulation polymers include cellulose acetate butyrate (CAB), ethyl cellulose (EC22 and EC100), low and medium molecular weight poly(methyl methacrylate) (PMML and PMMM), poly(alpha-methylstyrene) (PMS) and starch urea formaldehyde



(Starch-UF).

Examples of petroleum solvents include Aromatic 100, Aromatic 200, EXXSOL D 80, NORPAR 15, VARSOL 1, ISOPAR L, ISOPAR M, ISOPAR V and ORCHEX 796.

5 The pesticidal composition can be applied to plants in the form of sprays, dusts, or other formulations commonly employed in making pesticides. In the case of the plant material containing composition the material will be present in a dispersable or finely divided form to facilitate spraying onto plants to protect against pest attack. Such sprays would  
10 be useful in reducing pest numbers, whether the binding proteins, especially biotin-binding proteins, or degrading enzymes, had been released during processing via rupturing of the vacuoles, or not. If the vacuoles remain intact, then the proteins or enzymes will be released as the pests feed upon the preparation, and as such the invention may have utility as a mechanism for slow release of these proteins or enzymes, or any other proteins directed  
15 to the vacuole by the vector.

20 The compositions may further include one or more antifungal, antiviral, antimicrobial or pest control proteins all as discussed above. The use of these compositions in combination with the plants of the invention may be additive or synergistic, achieve broader spectrum control and reduce the risk of resistance developing.

25 Combinations particularly contemplated herein are compositions comprising proteinase inhibitors or insecticidal proteins such as *Bacillus thuringiensis* Cry proteins or biopesticides such as insect viruses or entomopathogenic fungi. Cry proteins including Cry1Ac, Cry1Cb, Cry1Da, Cry1F, Cry5 and Cry9A are preferred. The applicants have surprisingly found that plants transformed with biotin binding proteins and treated with Bt insecticidal protein exhibited synergistic toxic effects on pests (see Example 18). This suggests that plants containing chimeric genes expressing both biotin binding proteins and Bt proteins will be highly effective in protecting plants from pest attack. It is likely that such plants will  
30 be more toxic than those expressing either protein singly.

35 In another embodiment, the compositions of the invention can be used in conjunction with transgenic plants other than those of the invention. These other transgenic plants, for example, may incorporate genes conferring fungal, viral, microbial or herbicide resistance; genes conferring early ripening, heat stability, increased accumulation ability of desired products such as starch or cellulose or any other desirable trait as are known in the art. The composition of the invention when applied to the transgenic plant may also achieve the desirable results discussed above with plants of the invention.

40 In another embodiment, a composition of the invention may be applied to harvested material to prevent pest damage in storage. In an extrapolated application, the compositions may

similarly be used in plant derived products such as flours, meals, cereals and the like to prevent or control pest infestation.

Also provided by the present invention is a method for controlling or killing pests comprising administering to said pest an amount of a chimeric polypeptide of the invention, which includes a sequence encoding a pest control protein, effective to control or kill said pest.

In one embodiment of the method, the chimeric polypeptide is administered with a second pest control protein, wherein the combination provides more effective control than administration of the second pest control protein alone. It will also be appreciated that more complex combinations of pest control proteins including a polypeptide of the invention are feasible. Most commonly, two to five additional pest control proteins will be used. However, the methods and compositions are not limited thereto. The additional pest control plants may comprise any of those already discussed. A preferred additional pest control protein is a Bt protein, especially a Cry protein.

In a related aspect, also provided is a method of controlling or killing pests, the method comprising administering to said pest plant material of the invention which includes a sequence encoding a pest control protein. Compositions of the invention may also be used in these pest control methods.

It will also be appreciated that a further method for controlling pest attacks on plants of the invention expressing a pest control protein, comprises treating those plants with a Bt protein or composition incorporating same.

As discussed above, the pests against which the invention is most effective are the immature stages of insects, including larvae, grubs, nymphs and instars. Administration may be achieved according to any suitable method known in the art. For example, through plant material, sprays, mulches, baits, dusts or other compositions which the pest to be controlled takes up through feeding, inhalation, transdermal absorption or other administrative route. Pests which may be killed or controlled using this method include those discussed above and particularly those referenced in the accompanying Examples and those pests belonging to the same insect orders as those referenced in the accompanying Examples.

It will be appreciated that the above description is provided by way of example only and that variations in both the materials and techniques used which are known to those persons skilled in the art are contemplated.

Non-limiting examples illustrating the invention will now be provided.

**EXAMPLE 1****Experimental Details concerning the preparation of constructs****Materials:**

5 Custom primers were synthesized by Life Technologies. Subcloning Efficiency DH5 competent Cells were purchased from Life Technologies and the Hybaid Recovery Plasmid Mini Prep Kit from Hybaid Limited. All enzymes, unless otherwise stated were purchased from Promega. Ampligase Thermostable DNA Ligase and Reaction Buffer and GELase were purchased from Epicentre Technologies and Polymerase Chain Reaction (PCR)  
10 reagents from Perkin Elmer.

The Avidin cDNA (pGEMav) carried on the plasmid pGEM3 was supplied by Professor M. S. Kulomaa ((Department of Biological and Environmental Science, University of Jyvaskyla, Finland) and the Potato Proteinase Inhibitor I (PPI-I) cDNA was isolated in  
15 this laboratory (Beuning *et al.* 1994, GenBank Accession # L06606) and cloned into pUC19.

The Streptavidin cDNA, carried on the plasmid pET3a was supplied by The DuPont Merck Pharmaceutical Company. The Potato Proteinase Inhibitor II (PPI-II) genomic sequence was isolated in this laboratory and cloned into pUC19 (Murray and Christeller, 1994).  
20

**Methods:**

Subcloning Efficiency DH5 competent Cells were used for general cloning and amplification of recombinant plasmids and the Hybaid Recovery Plasmid Mini Prep Kit was used for plasmid preps. Isolation and recovery of DNA fragments was achieved by agarose gel  
25 electrophoresis followed by treatment of excised gel bands with GELase.

**DNA Sequencing and Computer Analysis:**

DNA sequencing was carried out on an Applied Biosystems (ABI) DNA Sequencer using  
30 dye terminator chemistry. Sequence analysis was performed using the Wisconsin Package Version 9.1, Genetics Computer Group (GCG), Madison, Wisconsin.

**EXAMPLE 2**

Preparation of a binary vector designed to express a chimeric polypeptide  
35 comprising Avidin mature peptide fused to a Potato Proteinase  
Inhibitor I Signal Peptide

**Methods:**

A one-step PCR-based mutagenesis method employing the combined use of a thermostable  
40 DNA polymerase and thermostable DNA ligase (Moore and Michael, 1995), was used to prepare a construct comprising the sequence encoding the mature Avidin polypeptide

- (Gope *et al.* 1987) fused to a PPI-I signal sequence. A Bgl II site was produced downstream of the PPI-I leader sequence at -positions 92 - 97 of the PPI-I coding sequence and a BamH I site was created upstream of the sequence encoding the mature Avidin polypeptide, at positions 65 - 70 of the sequence encoding the Avidin protein, as shown in Fig. 1 and Fig. 2 respectively. These two restriction sites have compatible cohesive ends.

**Primers:**

Forward M13 (lacZ) Primer [Perkin Elmer]:

10 5'-GCCAGGGTTTCCAGTCACGA-3'

Reverse M13 (lacZ) Primer [Perkin Elmer]:

5'-GAGCGGATAACAATTCACACAGG-3'

Avidin Upstream Primer:

5'-GCACACCCGGCTGTCCACCTG-3'

20 Phosphorylated Mutagenic Primers

PPI-I mutagenic primer:

5'-PGATGGACCAGAGATCTTAGAAC-3'

25 Avidin mutagenic primer:

5'-PGGCTCCCGGGATCCCTGCCAG-3'

**Amplification/Mutagenesis reactions:**

- To generate mutant products a total PCR reaction volume of 50  $\mu$ L with an effective 1 X Ampligase Reaction Buffer [20 mM Tris-HCl (pH 8.3 at 25°C), 25 mM KCl, 10 mM MgCl<sub>2</sub>, 0.5 mM NAD and 0.01% Triton X-100] was used with the following conditions:

- 100 pmol each outer primer  
1 nmol phosphorylated mutagenic primer  
40 nmol each dNTP  
0.1  $\mu$ mol dithiothreitol  
5 U Taq DNA polymerase  
5 U thermostable DNA ligase  
1ng recombinant plasmid DNA template

Reactions were first incubated at 94°C for 3 min., followed by 30 amplification cycles

performed as follows:

94°C, 1 min.

40°C, 1 min.

65°C, 6 min.

5

Amplification cycles were followed by a final extension at 65°C for 7 min.

Restriction analysis of amplification products from both mutagenesis reactions revealed mutant product to be present, but only at a maximum of 5% of the total product. To increase the yield of mutated product, Bgl II (for PPI-I mutagenesis) and BamH I (for Avidin mutagenesis) digestion products were ligated and then used as template for a second amplification reaction using outer primers only (Avidin Upstream and Reverse M13 (lacZ) for Avidin; Forward M13 (lacZ) and Reverse M13 (lacZ) for PPI-I). For PPI-I, greater than 95% of second round amplification product had the desired Bgl II site and approximately 80% of the second round product for Avidin mutagenesis possessed the BamH I site.

10  
15  
20  
25

The mutated PPI-I amplification product was digested with Bgl II and Sal I and the mutated Avidin product with BamH I and Hind III. The PPI-I leader sequence and the coding sequence for the Avidin mature protein were isolated and recovered for cloning along with Xho I/Hind III digested non-recombinant pART7 vector (Gleave, 1992). These three species were ligated, resulting in recombinant pART 7 [refer Fig. 5] and the sequence of the chimeric gene was checked. Subsequently, the expression cartridge containing the gene fusion was cloned into the Not I site of pART27 vector (Gleave, 1992) and this construct [refer Fig. 7A] was mobilized to *Agrobacterium tumefaciens* (strain LBA4404) by standard tri-parental mating techniques (Ditta *et al.* 1980).

### Discussion:

The resulting PPI-I/Avidin fusion protein has a total of 161 amino acids as shown in Fig. 8. The first 31 amino acids are PPI-I sequence and since the leader sequence comprises the first 23 amino acids, the original patterning of amino acids around with the site for cleavage between the signal sequence and the mature protein is retained. There are two single base pair changes in the gene fusion sequence relative to the predicted sequence. These changes are presumably the result of PCR error. One change is silent and the other results in an amino acid change from Serine to Proline at position 17 of the PPI-I signal sequence.

35

### EXAMPLE 3

Preparation of a binary vector designed to express a chimeric polypeptide comprising Synthetic "Core" Streptavidin peptide fused to a Potato Proteinase Inhibitor II Signal Peptide

40

**Methods:**

A fused gene was prepared comprising the sequence encoding Synthetic "Core" Streptavidin (Thompson and Weber 1993) fused to a PPI-II signal sequence. The Streptavidin cDNA, carried on the plasmid pET3a was cloned into the EcoR I/Xba I sites of pUC 19 (Fig. 3). The PPI-II signal sequence (Fig. 4) which contains an intron was isolated from recombinant plasmid using PCR with a sense primer binding to pUC19 and an antisense primer incorporating an EcoR I site into a 5' overhang. The primers were as follows:

sense primer:

5' - CTG CAG GTC GAC TCT AGA GGA - 3'

antisense primer:

5' - GGT GAA TTC TTA GTA CAG ATC TTC GCA - 3'

**Amplification reaction:**

A total PCR reaction volume of 50  $\mu$ l with an effective 1 X PCR Buffer [10 mM Tris-HCl, pH 8.3 and 50 mM KCL] was used with the following conditions:

20 pmol each primer  
15 nmol each dNTP  
2.0 mM MgCl<sub>2</sub>  
5 U Taq DNA polymerase  
1ng recombinant plasmid DNA template

Reactions were first incubated at 94°C for 2 min., followed by 30 amplification cycles performed as follows:

94°C, 1 min.  
50°C, 1 min.  
72°C, 1 min.

Amplification cycles were followed by a final extension at 72°C for 7 min.

The PCR product representing the PPI-II signal sequence was digested with Sal I and EcoR I. The recombinant plasmid pUC 19/Streptavidin cDNA was digested with EcoR I and Xba I and the Streptavidin cDNA was isolated from the vector and recovered. Non-recombinant pUC19 was digested with Sal I and Xba I and the three species were ligated to produce a construct comprising the gene fusion cloned into the Sal I and Xba I sites of pUC19. The sequence of the gene fusion was checked and subsequently cloned into the Xho I and BamH I sites of the pART7 vector [refer Fig. 6]. The pART7 expression

cartridge containing the gene fusion was then cloned into the Not I site of pART27 and this construct [refer Fig. 7B] was mobilized to *Agrobacterium tumefaciens* (strain LBA4404) by standard tri-parental mating techniques.

## 5 Discussion:

The resulting PPI-II/Streptavidin fusion protein has a total of 168 amino acids as shown in Fig. 9. The first 36 amino acids are PPI-II sequence. Five of these amino acids follow the cleavage site, preserving the amino acid pattern around this position. The nucleotide sequence of the PPI-II signal sequence includes a 119 bp intron (Murray and Christeller,  
10 1994).

## EXAMPLE 4

### Immunodetection of avidin in transgenic tobacco

#### 15 Methods:

##### 1. Tissue print

Samples were taken from the top 8 leaves of a tobacco plant expressing avidin (Pla2/9 #1). Four plants not expressing avidin were used as controls (PLA 2/3, NT12, GUS1  
20 and JB3-13).

Pieces of transgenic tobacco leaves 1 x 1cm were frozen at -20°C for 20 min, allowed to thaw and printed on to nitrocellulose using mechanical pressure.

#### 25 Labelling protocol:

Printed nitrocellulose membranes was washed in PBS-T (phosphate buffered saline with 0.1% Tween 20) for 20 min, blocked in 0.1% BSA-C (Aurion) for 15min and incubated in 1:1000 anti-avidin (Sigma A-5170) diluted in blocking buffer for 1h (as a control for non-specific binding, this last step was deleted in duplicate sets of prints). The membrane  
30 was then washed in PBS-T, incubated in goat anti-rabbit IgG-gold (10nm) (Sigma), washed again in PBS-T, then in double distilled water and drained. Finally the membrane was silver enhanced (BioCell silver enhancement kit) for 15min. Enhancement was stopped by washing in distilled water.

35

#### Results:

The nitrocellulose membrane silver enhanced (turned brown) over most of the tissue print area in the smallest top leaf. In all other leaves the silver enhancement was detected mainly towards the cut edges of leaf material. There was no silver enhancement on the prints  
40 from control plants or on the prints made in the absence of the anti-avidin antibody. This labelling protocol also acts as a test of the labelling procedure.

## 2. Embedded material

Pieces 1 x 1 x 5mm of transgenic tobacco leaf (Pla 2/9 #1) were fixed in 2% paraformaldehyde and 2.5% glutaraldehyde in 0.1M phosphate buffer under vacuum for 1h. The material was post-fixed in 1% osmium tetroxide 1h, dehydrated in an ethanol series and embedded in Spurr's resin. Pieces of non-transgenic tobacco (control material) were prepared in a similar manner. Sections were cut 300nm thick for light microscopy (LM) and mounted on Poly-L-lysine coated slides. Sections for electron microscopy (EM) were cut 130nm thick (gold) and mounted on carbon/formvar coated nickel grids.

## 10 **Labelling Protocol:**

For light microscopy the sections had a Pap pen ring drawn around them to contain the incubation liquid. The protocol for LM and EM were the same thereafter. The sections were etched for 30min in 10% hydrogen peroxide to remove the osmium, blocked in 0.1% BSA-c for 15 min, incubated in anti-avidin 1:500 in PBS-T for 1h (deleted for control) and washed in PBS-T. They were then incubated in goat anti-rabbit IgG-Alexa 488 (Molecular Probes) for 1h. The sections were then washed thoroughly in buffer and then in double distilled water.

The methodology for labelling of sections for the electron microscope (EM) was similar to that for the light microscope (LM) except goat anti-rabbit IgG-gold (10nm) was used instead of goat anti-rabbit IgG Alexa 488.

Sections were then viewed on a fluorescence microscope. Sections (1  $\mu$ m thick) were stained methylene blue/AzureII.

## **Results:**

Sections of Pla 2/9 #1 smallest top leaf stained for light microscopy shows darkly staining bodies mesophyll, epidermal cells, and cells of the glandular hairs (Figures 13 and 14)

Immunolabelling of LM (Figure 14) and EM (Figures 15 and 16) sections showed labelling of protein-type bodies in the vacuoles of mesophyll cells (both spongy and palisade) and in glandular hairs (Figures 15 and 16). The protein bodies were usually condensed into one body which was sometimes seen as a ring. There was no labelling in the vascular tissue or in the trichomes. Control material did not label.

## **Conclusions:**

The results indicate that avidin is synthesized in most common cell types in tobacco leaves. The bulk of the protein appears to be transported to the vacuole and deposited as a protein body within this organelle.



### EXAMPLE 5

#### ELISA assay of avidin and streptavidin

The following general ELISA assay technique was used for assaying for avidin and streptavidin where indicated in the following examples.

#### Method:

1. Plant material was ground with 2 volumes (w/v) of ice cold 0.05  $\mu$ M sodium phosphate (pH 7.5) containing 5% polyvinylpolypyrrolidone. This was centrifuged and the supernatant used for analysis. In order to construct standard curves control plant material was ground in the above buffer with and without 0.2 mg/mL avidin or streptavidin and centrifuged.

2. Generally 10  $\mu$ L of extract and 90  $\mu$ L of coating buffer (15 mM sodium carbonate, 46 mM sodium bicarbonate, pH 9.6) were mixed in a 96 well microtitre plate and incubated at 4C overnight. Each sample was duplicated and standards consisted of various proportion of control plant extract/added protein extract to the same total extract volume as the samples.

3. Plates were washed (3x) in phosphate-buffered saline (PBS) containing 0.02% Tween 20 (PBST) and incubated for 1 hr in 100  $\mu$ L of PBST containing 0.5% gelatin.

4. Plates were washed (3x) in PBST and incubated for 1 hr in 100  $\mu$ L of PBS containing either polyclonal rabbit anti-avidin or anti-streptavidin antibodies.

5. Plates were washed (3x) in PBST and incubated for 1 hr in 100  $\mu$ L of PBS containing goat anti-rabbit antibody linked to alkaline phosphatase.

6. Plates were washed (3x) in PBST and, after addition of 100  $\mu$ L of 0.1 M diethanolamine (pH 9.8) containing 0.5 mM  $MgCl_2$  and 0.5 mg/mL p-nitrophenyl phosphate, assayed at 410 nm in a microtitre plate reader. Initial rates of samples were determined by linear regression over 5-10 mins and compared to rates obtained for the duplicate standard curves (8 avidin or streptavidin concentrations) on each microtitre plate.

7. Concentrations of avidin and streptavidin in the samples were determined as the mean molar concentration in the tissue assuming that the specific gravity of plant tissue is one and molecular weights for avidin and streptavidin of 15600 and 16473 (for the standard) respectively.

### EXAMPLE 6

Toxicity of whole tobacco (*Nicotiana tabacum*) plants expressing avidin to potato

**tuber moth larvae (*Phthorimaea operculella*) (Lepidoptera: Gelechiidae)****Constructs:**

Non-transformed control plants

5 2 plants (NT 1, NT 2)

Control plants transformed with pumpkin fruit chymotrypsin inhibitor (PFCI) but not expressing the protein

3 plants (JB3/1, JB3/2, JB5/1)

Tobacco plants transformed with the avidin gene with a PPI-I targeting sequence (Example  
10 2 above)

6 lines (PLA2/2, PLA2/7, PLA2/9, PLA2/13, PLA2/20, PLA2/24), 4 clonal plants per line

**Trial Design:****Trial 1:**

The tobacco plants were removed from tissue culture and potted in fertilised potting mix (Smiths® general potting mix) before being placed in large ventilated acetate containers (220 x 300mm) in a containment glasshouse unit at  $22 \pm 5^\circ\text{C}$ . They were watered daily  
20 to maintain high humidity and soil moisture content.

Eight days later, when plants were well established with at least four to five small leaves, ten neonate potato tuber moth (PTM) larvae were placed on each tobacco plant, usually two per leaf. Prior to inoculation the larvae were weighed in batches of five (since single  
25 larvae are too small to give an accurate reading). TM larvae were obtained from a laboratory culture reared on potato tubers following the same basic procedure as Broodryk (1971) and Meisner et al. (1974).

**Trial 2:**

30 One week after Trial 1 was completed, the tobacco plants were cut back to the second node and allowed to regenerate leaves. When the plants had developed four to five leaves (in approximately 11 days) they were each inoculated again with ten neonate PTM larvae, usually two per/leaf, weighed in batches of five prior to inoculation as above.

**Trails 1 and 2:**

Inoculated plants were kept individually in acetate containers in the containment glasshouse unit at  $22 \pm 5^\circ\text{C}$  for nine days. Under these conditions growth of control larvae is exponential from hatch to nine days, but after this growth rate slows as pupation approaches. Hence in order to compare growth rates of larvae on control and transgenic plants, the trial was  
40 concluded after nine days. Damaged leaves containing larvae were removed, and larvae were dissected out of their mines within the leaf or stem tissue. The intention was to weigh

the larvae at this point in order to estimate growth rates, but, except for those on control plants, larvae were mostly dead, dried and shrivelled. Consequently head capsules were measured so that the instar reached at death could be recorded.

## 5 Level of expression of the avidin protein

### Results:

The level of expression of avidin in each of the plant lines was quantitated using chemiluminescence detection of avidin protein from western blots of leaf tissue, compared to authentic  
10 avidin standards and expressed as percentage of total leaf protein. These levels are given in Table 1 below.

**Table 1: The level of expression of avidin as % of total leaf protein, determined using the chemiluminescence method**

Plant Line	Avidin expression % total leaf protein ( $\mu$ M)*
PLA2/2	0.07 (0.90)
PLA2/7	0.10 (1.23)
PLA2/9	0.07 (0.90)
PLA2/13	0.06 (0.77)
PLA2/20	0.065 (0.83)
PLA2/24	0.06 (0.77)

\* The chemiluminescence method was used to estimate avidin expression as % total soluble  
25 leaf protein. In later Examples, an ELISA method (Example 5) was used to estimate the expression levels as  $\mu$ M. Hence these values were converted to  $\mu$ M. Avidin expression was measured in clones of these original plants using the ELISA method and results given  
30 in Example 8, Table 5. These levels are about three times higher than those given in Table 1. This may reflect the fact that, in these trials the measurements were done on plants still in tissue culture whereas those in Example 8 were done on large leaves from mature plants.

## 35 Mortality of PTM Larvae feeding on whole tobacco plants expressing the avidin gene.

### Trial 1:

Good recovery rates of larvae from both control and transgenic plants were obtained:  
86% from controls and 76.7% from transformed plants. Fig. 10 clearly shows the high  
40 mortality PTM larvae after feeding for nine days on whole transgenic tobacco plants expressing

the avidin gene compared to both non-transformed control plants and control plants transformed with, but not expressing, the pumpkin fruit chymotrypsin inhibitor (PFCI) gene.

5 The majority of dead larvae were recovered from mines where they had died at the "cutting face". A few (5% of dead larvae) were recovered from the surface of leaves, having generally left a mine close by. It is most likely that the majority of larvae not recovered had died in this way and had fallen off the leaves. Some mines were found without occupants. However, there was no evidence that larvae had started and abandoned mines on several occasions as we have previously observed in another experiment in which larvae were  
10 fed on tobacco expressing *cry* 1Ac and *cry* 9Aa2 genes (Gleave et al. 1998).

PTM larvae undergo four instars during their development. In order to define the stage of development of the larvae at death, head capsule widths were measured using a micrometer eye-piece. All control larvae were alive and most were third instars. None of the larvae recovered on any of the plants expressing avidin had reached third instar before death and many had died during or just after the moult from first to second instar, as evidenced by the fact that the ecdysed cuticle was still attached. This reflects results in earlier trials with avidin incorporated into diet. Table 2 below gives a breakdown of instars on each plant line.

**Table 2: Number of larvae at each instar recovered from transgenic tobacco plants expressing avidin in Trial 1**

	Plant line	Neonates inoculated	Number of larvae at			
			1st instar	2nd instar	3rd instar	4th instar
25	NT control	20	0	1	18	0
	JB control	30	0	0	23	1
	PLA2/2	40	3	28	0	0
	PLA2/7	40	4	23	0	0
	PLA2/9	40	2	27	0	0
30	PLA2/13	40	1	25	0	0
	PLA2/20	40	2	27	0	0
	PLA2/24	40	4	25	0	0

#### **Trial 2:**

35 Again there were good recovery rates of larvae from both control and transgenic plants: 88% from controls and 88.8% from transformed plants. Fig. 11 reflects the results of the first trial showing high mortality of PTM larvae fed on whole transgenic tobacco plants expressing the avidin gene compared to those on control plants. In fact a total of only four live larvae were recovered from all avidin expressing plants (<1.7% survival), whereas  
40 only three larvae had died on the control plants (94% survival).

Head capsule widths of larvae were measured and the number of recovered larvae at each instar is given in Table 3. As in the first trial, none of the larvae recovered from any of the plants expressing avidin had reached third instar before death and many had died during or just after the moult from first to second instar; again the ecdysed cuticle was still attached in many cases.

**Table 3: Number of larvae at each instar recovered from transgenic tobacco plants expressing avidin in Trial 2**

Plant line	Neonates inoculated	1st instar	Number of larvae at		
			2nd instar	3rd instar	4th instar
NT control	20	0	3	12	0
JB control	30	0	1	24	3
PLA2/2	40	34	5	0	0
PLA2/7	40	25	9	0	0
PLA2/9	40	30	3	0	0
PLA2/13	40	26	11	0	0
PLA2/20	40	25	5	0	0
PLA2/24	40	30	3	0	0

#### Conclusion:

Total mortality of PTM larvae fed on tobacco plants expressing the avidin gene would have occurred if the trials had been continued beyond nine days; larvae that survived for nine days were small, shrivelled and close to death as evidenced by their minimal response when touched by a fine sable paint brush.

Avidin expressed in tobacco plants is highly toxic to PTM larvae and has definite potential in the development of pest resistant crop cultivars.

#### EXAMPLE 7

**Toxicity of whole tobacco (*Nicotiana tabacum*) plants expressing streptavidin to potato tuber moth larvae (*Phthorimaea operculella*) (Lepidoptera: Gelechiidae)**

#### Constructs:

Non-transformed control plants

6 plants (NT21-26)

Plants transformed with and expressing the streptavidin gene with a PPI-II targeting sequence (Example 3 above) - (Sav)

6 plant lines, 5 clones per line (5, 9, 10, 14, 23, 26)

2 plant lines, 3 clones per line (25, 28).

**Trial Design:****Trial 1:**

5 The transformed tobacco plants were removed from tissue culture, planted in fertilised potting mix (Smiths® general potting mix) and placed individually in large ventilated acetate containers (220 x 300mm) in a containment glasshouse unit at  $24 \pm 7^\circ\text{C}$ . They were watered regularly to maintain high humidity and soil moisture content until well established.

10 Eleven days later, five neonate potato tuber moth (PTM) larvae were placed on each tobacco plant. Prior to inoculation the larvae were weighed in batches of five (since single larvae are too small to give an accurate reading on a 5-place balance). PTM larvae were obtained from a laboratory culture reared on potato tubers following the same basic procedure as Broodryk (1971) and Meisner et al. (1974).

**Trial 2:**

15 On completion of Trial 1, the tobacco plants were cut back to the second node and allowed to regenerate new leaves. When the plants had developed at least four to five leaves they were each inoculated again with five neonate PTM larvae as above. Unfortunately some individual plants died during this process and so fewer clones were tested for some lines in the second trial.

**Trials 1 and 2:**

25 Inoculated plants were kept individually in acetate containers in the containment glasshouse unit at  $24 \pm 7^\circ\text{C}$  for nine days. Under these conditions growth of control larvae is exponential from hatch to nine days, but after this growth rate slows as pupation approaches. Hence in order to compare growth rates of larvae on control and transgenic plants, the trial was concluded after nine days. Damaged leaves containing larvae were removed, and larvae were dissected out of their mines within the leaf or stem tissue. The intention was to weigh  
30 the larvae at this point in order to estimate growth rates, but, except for those on control plants, the larvae were mostly dead, dried and shrivelled. Consequently head capsule width was measured for all larvae retrieved so that the instar reached at death could be recorded.

**Results:****Level of expression of the streptavidin protein:**

The level of expression of streptavidin in each of the plant lines was measured using the technique described in Example 5. These levels are given in Table 4 below.

Table 4: Expression of streptavidin in tobacco plants

Plant Line (Savα)	Expression of Streptavidin μM (s.e.)
5	12.802 (0.834)
9	17.818 (0.059)
10	11.404 (0.896)
14	18.178 (0.560)
23	24.524 (0.042)
25	21.703 (0.842)
26	16.306 (1.831)
28	15.788 (0.260)

Mortality of PTM larvae feeding on whole tobacco plants expressing the streptavidin gene:

#### Trials 1 and 2:

Recovery of larvae was good from control plants in both trials (88.6 and 92% respectively) and from transgenic plants (78.3 and 83% respectively) and similar to that reported in the trials with tobacco expressing the avidin gene (Example 6). Figure 17 shows the number of live and dead larvae recovered nine days after inoculation, from each plant line in both trials. In Trial 2 there was total mortality on all plant lines, but in Trial 1 there were a few survivors after nine days on some plant lines: of the 25 larvae initially placed on the plants, two survived on line 5 and one each on lines 9, 10, 23 and 28. However, all of these "survivors" were close to death. In contrast, there was no larval mortality on control plants in either trial. As in Example 6, the majority of larvae had died within the mines in the leaves and only a few dead larvae were found on the leaf surface after abandoning their mines.

Head capsule widths of all larvae were measured after they were removed from their leaf mines to determine their stage of development. Larvae recovered from non-transgenic (NT) plants were all alive in both trials and all but two had reached 3rd or 4<sup>th</sup> instar. In contrast, the majority of larvae feeding on the transgenic plants had died at 1<sup>st</sup> or 2<sup>nd</sup> instar (Figure 18). Most of these had died just prior to or during the ecdysis from 1<sup>st</sup> to 2<sup>nd</sup> instar as was evidenced by the number of dead larvae with ecdysed skins and head capsules still attached.

#### Conclusions:

Tobacco plants expressing the streptavidin gene were highly insecticidal to potato tuber moth larvae. Larval mortality occurred on all plants tested expressing the gene and the majority of larvae died just prior to, during, or immediately after ecdysis between the 1<sup>st</sup> and 2<sup>nd</sup> instar.

**EXAMPLE 8**

Toxicity of avidin expressed in tobacco (*Nicotiana tabacum*) leaves to larvae of the common cutworm *Spodoptera litura* (Lepidoptera: Noctuidae) and the cotton bollworm (tomato fruitworm, cornear worm) *Helicoverpa armigera* (Lepidoptera: Noctuidae)

**Constructs:****Control lines:**

10 Non-transformed control plants:

4 plants (NT11, NT12, NT13, NT14)

Control plants transformed with PRD400 vector with pumpkin fruit chymotrypsin inhibitor (PFCI) gene but which do not express the transgene:

15 8 plants (6 independent transformants) (JB3-1C/AB, JB3-1, JB3-13, JB3-15, 2 clonal JB3-16 plants, and 2 clonal JB3-25 plants)

**Control plants transformed with the pART27 vector:**

20 7 plants (all independent transformants) (art27c #1, art27c #3, art27c #4, art27c #5, art27c #6, art27c #7, art27c #8)

**Control plants containing the pART27 vector with the GUS gene (uid):**

4 plants (all independent transformants) (GUS1, GUS2, GUS5, GUS8)

**25 Avidin-expressing lines:**

Tobacco plants transformed with the avidin gene with a PPI-I targeting sequence (Example 2 above):

6 plant lines derived from 6 separate transformation events (PLA2/2, PLA2/7 PLA2/9, PLA2/13, PLA2/20, PLA2/24), 4 clonal plants per line.

30

**Insects:**

*Spodoptera litura* were obtained from a laboratory colony originally established from moths field-collected in Queensland, Australia, while *Helicoverpa armigera* were from a laboratory colony established from moths collected in Christchurch, New Zealand.

35 Both colonies were reared on artificial diet as described in McManus and Burgess (1995).

Neonate *S. litura* larvae were placed on tobacco leaves within 12h of emergence from



eggs. Initial larval weight was estimated from the mean weight of three samples of 100 larvae.

- 5 Neonate *H. armigera* larvae were placed on artificial diet for 48h following emergence from eggs, and then placed on tobacco leaves as late first instar larvae. Initial larval weight was determined as the mean of the individual weights of a randomly chosen sample of 48 larvae weighed at the beginning of the experiment.

#### Trial design:

- 10 On each plant used in the experiment, Leaf 1 was designated as the uppermost (youngest) leaf which was 15cm or more in length from leaf tip to leaf base (the point at which the leaf joined the petiole). The leaves below Leaf 1 were assigned numbers consecutively down the plant. Leaves 1 and 2 were used for *H. armigera* as previous experiments had shown that larvae grow best on these leaves. For similar reasons, *S. litura* were given  
15 Leaves 4 and 5 of the same plants.

To ensure leaves remained turgid during larval feeding, each leaf was cut from the plant close to the stem leaving a long petiole, and each petiole was immediately plunged into about 20 mL of a setting solution of 0.4% agar in a 30 mL coulter cup.

- 20 At the start of the experiment, larvae were placed on leaves from one plant from each of the six clonal avidin-expressing lines, and on six control plants. Twelve *H. armigera* larvae were placed on the undersides of Leaves 1 and 2, i.e. 12 larvae x 2 leaves x 6 plant lines = 144 larvae, and an equivalent number of control larvae were used. For *S. litura*,  
25 15 larvae were placed on the upper surfaces of Leaves 4 and 5, i.e. 15 larvae x 2 leaves x 6 plant lines = 180 larvae on both avidin and control treatments.

- Each leaf with larvae was placed in a 300 x 210 x 80mm plastic storage box lined with paper towels and with a snap-on lid. Larvae and leaves were checked daily, and leaves  
30 were replaced with new ones from fresh plants as necessary so that larvae could feed *ad libitum*. Throughout the experiment, larvae on avidin plants were fed leaves from within the same clonal line (e.g. PLA2/2 or PLA2/7), and larvae on control plants were kept on the same genetic plant type (NT or JB-3 or art27c or GUS). When necessary, leaves of the equivalent physiological age from previously used plants were utilised.

- 35 The experiment was conducted in a controlled temperature room at  $24.5 \pm 1^\circ\text{C}$  and 60% relative humidity, with a 16:8h light:dark cycle.

Larvae were first weighed and survivors counted at Day 8, and then at regular intervals

throughout the experiment until death or until pupation had begun in a treatment.

#### Determination of avidin expression levels:

To measure expression levels in plants fed to larvae, two leaf samples of 8cm<sup>2</sup> were taken from Leaf 4 of all avidin plants used in the trial. One sample was taken just before larvae were initially placed on the leaf, and the other a few days later, following the transfer of larvae from the leaf onto a fresh leaf. Expression was measured as described in Example 5.

Table 5: Expression levels of avidin in plants

Plant line	Mean expression level of avidin ( $\mu$ M)	Standard error	Number of samples
PLA2/2	3.10	0.42	8
PLA2/7	3.29	0.23	8
PLA2/9	4.37	0.51	8
PLA2/13	3.40	0.38	8
PLA2/20	4.59	0.31	8
PLA2/24	4.10	0.21	8
NT	0	-	8

#### Results:

##### *S. litura*

As the same pattern of response was observed in larvae on all control lines, results from the different lines were pooled. The same observation was made for larvae on all avidin lines, so results from these lines were also pooled.

Larvae on avidin-expressing plants were significantly smaller than controls at the first weighing on Day 8 (control plants: N = 153, mean weight = 0.0304 g, s.e. = 0.0014; avidin plants: N = 160, mean weight = 0.0151 g, s.e. = 0.0007;  $P < 0.001$ ) (Fig. 19), (ANOVA, Payne *et al.*, 1993), but there were no differences in survival at that time. By Day 12, larvae eating avidin plants had begun to die ( $P < 0.001$ ) (Fig. 20), and there were clear differences in mean weight and total live biomass present on the two treatments ( $P < 0.001$ ) (Fig. 21). By Day 15, these differences were even more pronounced, and after this time control larvae had pupated, so no further control measurements were taken. Comparative larval sizes on control and avidin plants are shown on Day 15 in Figure 19B. Differences in size and plant damage on Day 15 are shown in Figure 19C. Larvae on avidin plants steadily diminished in numbers and total biomass, and by day 25 all had died.

We observed that larvae feeding on avidin plants were unable to successfully complete the process of moulting from one instar to the next. Larvae on these plants appeared to stop feeding during ecdysis, and to then turn black and die while still attached to a partially shed larval skin.

### *H. armigera*

As with *S. litura*, larval responses on all control lines were the same, and results were thus pooled, as were responses on all avidin lines. *H. armigera* larvae fed avidin-expressing plants were smaller than those fed control plants by Day 8 (control plants: N = 130, mean weight = 0.0909 g, s.e. = 0.0031; avidin plants: N = 130, mean weight = 0.0375 g, s.e. = 0.0013;  $P < 0.001$ ) (Fig. 22). Three days later, control larvae had continued to grow well, while avidin-fed larvae had begun to die ( $P < 0.001$ ) (Fig. 23), and differences in biomass between the two treatments were extreme ( $P < 0.001$ ) (Fig. 24). No further control measurements were made after Day 11 as larvae had begun to pupate. Comparative larval sizes on control and avidin are shown on Day 14 in Figure 22 B. Differences in size and plant damage on Day 14 are shown in Figure 22C. By Day 22, all larvae on avidin plants had died.

As with *S. litura*, *H. armigera* larvae on avidin plants often died during the moulting process.

### Conclusions:

The expression of avidin in six different transgenic lines of tobacco was fatal to larval *S. litura* and *H. armigera*. Larvae of both these lepidopteran pest species grew rapidly and pupated on a range of non-transgenic and transgenic tobacco lines which did not express avidin. Larvae fed avidin-expressing plants were unable to develop normally or attain significant biomass, often dying during early instar moults.

These results provide further evidence of the effectiveness of the avidin construct described in this patent in protecting the plant in which it is expressed from insect damage.

**EXAMPLE 9**

**Toxicity of avidin expressed at a range of concentration levels in tobacco (*Nicotiana tabacum*) leaves to larvae of the cotton boll worm (tomato fruitworm, cornear worm) *Helicoverpa armigera* (Lepidoptera: Noctuidae)**

5

**Constructs:****Control lines:****Non-transformed control plants:**

10 48 plants (NT 101 - NT 148).

These were grown from seeds produced by selfed NT plants 11-14 which were used in the trial described in Example 8.

15 **Avidin-expressing lines:**

Tobacco plants ( $T_1$ ) were grown from seeds collected from 3 selfed plants from the clonal lines PLA2/7, PLA2/9 and PLA2/13. These  $T_0$  parental plants had been transformed with the avidin gene with a PPI-I targeting sequence (Example 2 above), and were used in the trial described in Example 8. Twenty four plants from each of these three  $T_1$  seed lines were grown for the experiment, and 25 of these 72 plants were selected for use depending on their level of avidin expression.

20 **Insects:**

Neonate *H. armigera* larvae from our laboratory colony (see Example 8) were placed on the leguminous host plant *Lotus corniculatus* and kept at 18°C for 3-4 days prior to the experiment. Late first instar larvae were then transferred to control and avidin-expressing tobacco leaves. Initial larval weight was determined as the mean of the individual weights of a randomly selected sample of 48 larvae weighed at the beginning of the experiment.

30 **Trial design:**

To measure the effect of avidin expression level on the growth, survival and biomass of *H. armigera* larvae, the 72  $T_1$  avidin plants described above were tested for expression level using the ELISA method (Example 5). A leaf sample of approximately 50-60cm<sup>2</sup> was removed from the tip of Leaf 4 of each plant for this process. All plants were ranked according to their expression level, and divided into six groups representing six non-overlapping ranges of expression level, from "high" to "low". These six groups of plants were assigned as six treatments with different mean avidin concentrations (Table 6).

At the start of the trial, the highest expressing plant from each treatment group and two

control plants of similar physiological form were selected. Leaf numbers were assigned on each plant as described in Example 8, and Leaves 1 and 2 cut from each plant for use in the trial. Petioles were again immersed in setting in agar to maintain leaf freshness, and 12 *H. armigera* larvae were placed on the underside of each leaf.

5

As in Example 8, leaves were stored in plastic boxes, and the experiment conducted at  $24.5 \pm 1^\circ\text{C}$  and 60% relative humidity, with a 16:8h light:dark cycle.

10 Larvae were weighed on Days 8, 11, 13, 14 and 15, and surviving larvae were transferred to fresh leaves 1 and 2 from the next highest expressing plant in each treatment group on Days 6, 8, 11 and 16. To ensure that larvae could feed *ad libitum*, additional leaves were cut from positions immediately above or below Leaves 1 and 2 on the same plants, and provided to larvae if necessary. Control larvae required many more leaf additions than all other treatments, and thus were given additional leaves from a range of control  
15 plants and leaf positions.

Table 6: Expression levels of avidin in treatment groups of plants used in trial

Treatment	Mean expression level of avidin ( $\mu\text{M}$ )	Standard error	Number of plants used
1	17.25	0.44	5
2	14.18	0.09	4
3	10.85	0.07	4
4	8.71	0.12	4
5	6.40	0.12	4
6	3.69	0.11	5
Control	0	-	48

#### - Results:

30 As there were no significant differences between larval growth, survival and biomass on the two control treatments, the results of these two treatments were combined.

By the time larvae were first weighed on Day 8 of the experiment, control larvae had grown larger than those in all other treatments (Fig. 25) ( $P < 0.05$  -  $P < 0.0001$ ) (ANOVA., Payne *et al*, 1993). These differences increased with time.

35

Comparison of larval survival curves using a log-rank test (Kalbfleisch and Prentice, 1980) showed that survival on all avidin-expressing lines was significantly reduced in comparison with control survival ( $P < 0.001$ ). There were no significant differences between survival on any of the six lines expressing avidin at different levels ( $P = 0.328$ ).

All the larvae fed plants expressing avidin at 17.25 - 6.40  $\mu$ M failed to achieve substantial growth, and died, often during moulting, without pupating (Fig. 26). Two of the 24 larvae on the lowest expressing avidin treatment pupated, although they were smaller than control larvae. One of these pupae emerged as a moth. On the two control treatments, 31 of 48 larvae successfully pupated and 19 of these emerged as moths. The number of larvae successfully pupating in the control treatments was reduced by cannibalism of prepupae by voracious late instar larvae. This effect may also have reduced the rate of emergence of moths from pupae in the controls. No such effect occurred in the avidin treatments because of the extremely high larval death rate caused by the ingestion of avidin-expressing leaf material.

Accumulation of biomass on the avidin-expressing lines was negligible compared to that on the control lines (Fig. 27).

#### Conclusions:

Tobacco plants expressing avidin at levels ranging from 6.40 to 17.25  $\mu$ M caused total mortality of *H. armigera* larvae in this trial. Expression levels of 3.69  $\mu$ M resulted in a very high level of larval mortality (92%). All plants expressing avidin at any level were protected from insect attack as evidenced by the extremely low biomass of insects on those plants.

### EXAMPLE 10

**Toxicity of avidin and streptavidin incorporated into artificial diets to the pine shoot tip moth, *Rhyacionia buoliana* (Lepidoptera: Tortricidae)**

#### Insects:

A laboratory colony of *Rhyacionia buoliana* was established by field collection of late instar larvae and pupae from pine (*Pinus radiata*) plantations throughout Chile. Field-collected individuals which became adults were confined in laboratory cages to allow mating. Eggs laid by adult females were collected, and larvae which emerged from these were used in this trial.

#### Methods:

The avidin used in this experiment was a Calbiochem® product, purchased from Calbiochem-Novabiochem Corporation, La Jolla, CA 92039. It was lyophilized avidin from egg white. Product Number 189725, Lot Number 276992.

The streptavidin was also obtained from Calbiochem-Novabiochem Corporation, and was a lyophilized solid, Product Number 189730, Lot Number B19870.

Avidin and streptavidin were incorporated into artificial diet at the following concentrations

5 in eight treatments:

1. control, 0  $\mu\text{g}/\text{mL}$
2. control, 0  $\mu\text{g}/\text{mL}$
3. avidin, 50  $\mu\text{g}/\text{mL}$
4. avidin, 100  $\mu\text{g}/\text{mL}$
- 10 5. avidin, 1000  $\mu\text{g}/\text{mL}$
6. streptavidin, 50  $\mu\text{g}/\text{mL}$
7. streptavidin, 100  $\mu\text{g}/\text{mL}$
8. streptavidin, 1000  $\mu\text{g}/\text{mL}$

15 These levels are equivalent to plant expression of 3.2, 6.4 and 64  $\mu\text{M}$  of avidin, and 3.0, 6.1 and 60.6  $\mu\text{M}$  of streptavidin. We have shown avidin expression levels in tobacco ranging from 3-25  $\mu\text{M}$  (Examples 8, 9 and 18), and streptavidin levels of 11-24  $\mu\text{M}$  (Example 7).

20 The artificial diet used in this experiment was a general purpose insect rearing diet based on the recipe of Singh (1983). The avidin and streptavidin were added in aqueous solution into freshly made diet, which had cooled to 60°C.

25 The experiment was run in a randomised complete block design, in three blocks, which were set up on consecutive days. Both avidin and streptavidin, at each of the three doses, were fed to a total of 90 larvae, and 180 larvae were given control diet:

i.e. 2 proteins x 3 concentrations x 30 larvae x 3 blocks = 540 larvae  
+ 2 controls x 30 larvae x 3 blocks = 180 larvae

30

Within 12h of hatching from eggs, neonate larvae were placed in pottles containing BIO-SERV® pine tip moth diet (the diet on which the colony was reared).

35 At the beginning of the experiment, healthy 24h-old larvae which had established well on this diet were then transferred to 1.5 mL Eppendorf tubes containing a 0.25 mL block of treatment diet, where they were confined individually. Initial mean larval weight was determined by weighing 100 of these healthy larvae selected for the experiment *en masse*.

Larval survival was checked every seven days for the duration of the experiment. After 14 days, larvae were weighed and transferred to new tubes with 1 mL of fresh diet. After 35 days, surviving larvae were weighed again, and the experiment terminated.

- 5 This experiment was conducted in a temperature-controlled incubator set at 20°C, in which lights periodically switched on when the temperature dropped below the target.

### Results:

- 10 As there were no significant differences between data collected for the three blocks of any given treatment, results for the three blocks were pooled within all treatments.

Both avidin and streptavidin at all 3 concentrations had caused significant reductions in larval growth by Day 14 (Fig. 28) ( $P < 0.0001$ ) (ANOVA, Payne *et al.*, 1993), and these differences increased by Day 35. Both proteins were toxic to larvae, and most individuals  
15 feeding on an avidin or a streptavidin diet were dead before the end of the experiment (Fig. 29). Many of the dead larvae had died during the process of moulting from one instar to the next. Larvae that survived feeding on diet containing either protein at any of the three concentrations were close to death. Comparison of survival curves using  
20 log-rank tests showed all treatments reduced larval survival compared with controls ( $P < 0.001$ ). The highest dose of streptavidin killed larvae faster than any of the other treatments ( $P < 0.001$ ), but there were no other differences among the survival responses to other doses of either protein. Because both avidin and streptavidin killed most larvae and prevented growth in survivors, there were very large differences between insect biomass on controls and all other treatments (Fig. 30).

25

### Conclusions:

- This trial has demonstrated the high level of toxicity of both avidin and streptavidin to the pine shoot tip moth, *Rhyacionia buoliana*. These results suggest that either of these proteins would control the pest if expressed in *P. radiata* or other host trees at levels equivalent  
30 to those we have demonstrated for avidin and streptavidin in tobacco plants (Examples 7, 8, 9 and 18).

### EXAMPLE 11

- 35 Toxicity of avidin-painted willow (*Salix fragilis*) leaves to neonate willow sawfly larvae (*Nematus oligospilus*) (Hymenoptera: Tenthredinidae).

### Insects:

Willow sawfly larvae (*Nematus oligospilus*) which had hatched within the previous 24-hour period, were obtained from a laboratory colony reared on small potted willow plants (*Salix*



*fragilis*).

**Leaf material:**

- Leaves were obtained from potted willow plants (*S. fragilis*) grown in a shade house, the same source as those on which the larvae were reared.

**Methods:**

- The avidin used in this trial was a Calbiochem® product, purchased from Calbiochem-Novabiochem Corporation, La Jolla, CA 92039. It was lyophilized avidin from egg white, Lot 276992.

Willow leaves were weighed and a mean leaf weight obtained ( $194.5 \pm 13.1$  mg). Using this weight the amount of avidin to apply per leaf was calculated as 65 and 130  $\mu$ M delivered as 200 and 400  $\mu$ g avidin/leaf.

- To ensure avidin was well distributed over the leaf surfaces it was dissolved in a 0.1% solution of the "wetter and sticker", BondXtra® (i.e. 50  $\mu$ L in 50 mL). 100  $\mu$ L /leaf gave good coverage.

- Solutions were painted on to leaves using a sable brush (Cirrus 110®). The brush was weighed before and after applying the solutions to the leaves and was found to absorb about one-tenth of the volume. Hence 55  $\mu$ L of each solution was pipetted on and applied to each side of each willow leaf. Leaves were allowed to air dry.

**Trial design:**

- Excised leaves were trimmed to fit across a Petri dish, one leaf per dish. The leaf petiole was placed in a small tube of water and painted with the appropriate solution. After being air-dried, the leaf was then pushed through a hole in the side of the Petri dish. Water was topped up every 2 days. Close cell foam supported the petiole and filled the space around the hole preventing the larvae escaping. The Petri dish with tube attached was stuck to a backing board with Blu-tack® and held firmly in place with a rubber band. The whole set up was then set vertically on a slotted board. Each Petri dish contained one willow leaf and one larva and each treatment tested 20 larvae. There were four treatments:
1. controls in which leaves were untreated,
  2. 0.1% BondXtra®,
  3. 65  $\mu$ M avidin in 0.1% BondXtra®,
  4. 130  $\mu$ M avidin in 0.1% BondXtra®.

Larvae were weighed individually and one placed in each Petri dish containing a single willow leaf. Surviving larvae were weighed again after 7, 14 and 21 days and leaves were changed after 10 and 15 days.

## 5 Results:

Figure 31 shows survival of sawfly larvae over the first 21 days by which time the majority of survivors had pupated. Whilst no controls died and only one death was recorded amongst larvae treated with BondXtra<sup>®</sup> survival of larvae on leaves coated with avidin declined steadily. The proportion of sawfly larvae surviving to 21 days on leaves coated with 65  $\mu$ M avidin was 0.4, and 130  $\mu$ M avidin only 0.1. Further, weight gain over the first 14 days was significantly reduced at both avidin concentrations when compared to control larvae and those feeding on leaves treated with BondXtra<sup>®</sup> alone (Figure 32).

At pupation, sawfly larvae form a fibrous pupal case or cocoon. At the lower avidin concentration only one out of the six larvae that reached pupation and developed a cocoon failed to emerge as an adult. At the higher avidin concentration, only one larvae attempted and failed to pupate; no adults emerged from this treatment (Figures 33 and 34). In both cases where the larva failed to emerge as an adult the fibrous pupal case contained a shrivelled dead larval body and so ecdysis (moult) had not been completed.

## 20 Conclusions:

Avidin is highly insecticidal to willow sawfly larvae and, as has been observed in bioassays with this protein on other insect species (see other examples), it appears to have acted both as a growth inhibitor and as a moulting inhibitor.

## 25 EXAMPLE 12

Toxicity of avidin-painted lettuce (*Latuca sativa*) leaves to the black field cricket, *Teleogryllus commodus* (Orthoptera: Gryllidae)

## 30 Insects:

Crickets were obtained from a laboratory colony of *Teleogryllus commodus* originally field collected in Northland, New Zealand. Four day old nymphs were used in this trial. These individuals had been fed since eclosion from eggs on the normal colony diet for young nymphs of rolled oats, dried lucerne (*Medicago sativa*) meal and dog biscuits (Pedigree<sup>®</sup>

35 PAL Meaty-Bites<sup>®</sup>).

## Methods:

The avidin used in this trial was a Calbiochem<sup>®</sup> product, purchased from Calbiochem-Novabiochem Corporation, La Jolla, CA 92039. It was lyophilized avidin from egg white,

Lot 276992.

Green distal portions of leaves from organically grown lettuce leaf were cut into sections approximately 4 x 4 cm. These were painted on both sides with three different solutions, providing three treatments:

1. Control solution of 0.1% (v:v) BondXtra<sup>®</sup>, a wetting, spreading and sticking agent
2. 4.8  $\mu$ M avidin (75  $\mu$ g/g fresh weight of lettuce leaf) in 0.1% (v:v) BondXtra<sup>®</sup>
3. 19.2  $\mu$ M avidin (300  $\mu$ g/g fresh weight of lettuce leaf) in 0.1% (v:v) BondXtra<sup>®</sup>

- Cricket nymphs were weighed and placed individually in 75 mL specimen pottles with ventilation holes punched in their lids, and with a 42.5 mm filter paper disc placed in the bottom of each pottle to absorb excess moisture. Food was replaced as necessary so that crickets could feed *ad libitum* on fresh leaf material. Each cricket was weighed weekly until all individuals feeding on the avidin-painted leaves had died.

#### Results:

- Cricket nymphs grew well on control leaves but poorly on leaves painted with avidin at both concentrations (Figure 35). By Day 21 control crickets were significantly larger than those surviving avidin treatment ( $P < 0.05$ ) (ANOVA, Payne *et al.* 1993). By this time, all those on 4.8  $\mu$ M avidin leaves were dead (Figure 36) and there were few survivors on the 19.2  $\mu$ M avidin treatment. By Day 35, all crickets on the 19.2  $\mu$ M treatment had also died. Many of the avidin-fed crickets died while moulting from one nymphal stage to the next. Cricket biomass in the control treatment steadily increased throughout the experiment, while biomass had reached zero in the 4.8  $\mu$ M avidin treatment by Day 21, and dropped below the starting value in the 19.2  $\mu$ M treatment by this time (Figure 37). Biomass in the 19.2  $\mu$ M treatment fell to zero soon after this.

#### Conclusions:

- Avidin is highly toxic to the black field cricket, demonstrating the efficacy of this protein as a means of controlling orthopteran pests. This suggests the use of avidin-expressing plants as a means of controlling pests such as locusts and grasshoppers as well as crickets.

#### EXAMPLE 13

- Toxicity of artificial diet containing streptavidin to neonate clover root weevil (*Sitona lepidus*) (Coleoptera: Curculionidae) and neonate Argentine stem weevil (*Listronotus bonariensis*) (Coleoptera: Curculionidae)

#### Insects:

Eggs of both weevil species were obtained from field-collected adults maintained on white

clover, *Trifolium repens*, (for *Sitona lepidus*) and ryegrass, *Lolium perenne*, (for *Listronotus bonariensis*) foliage.

5 *S. lepidus* eggs were placed in Petri dishes on filter paper moistened with sterile distilled water and allowed to hatch at 25°C. To delay hatching until sufficient eggs had been laid for a trial, some eggs were stored for up to 24 days at 10°C, before being brought to the higher temperature for hatching.

10 *L. bonariensis* eggs were placed directly onto blocks of artificial diet in small plastic containers (4 mL autoanalyser cups), one larva per cup.

### Streptavidin:

15 The streptavidin used in this trial was obtained from Calbiochem-Novabiochem Corporation, and was a lyophilized solid. Product Number 189730, Lot Number B19870.

### Diets:

20 An artificial diet (ASW diet) known to be suitable for rearing *L. bonariensis* (Malone and Wigley, 1990) was modified by omitting biotin from the recipe and used in the streptavidin trials for both weevil species.

25 To test the diet's suitability for *S. lepidus* trials, some of the first neonate *S. lepidus* larvae obtained were placed onto blocks of unmodified ASW diet (with biotin) for three days prior to being used in the first replicate of the streptavidin feeding trial. Other larvae in this trial had been maintained for the first three days of life on either washed clover roots or a second artificial diet, which also contained biotin (porina diet) (Burgess *et al.*, 1993). As larval feeding was observed only on ASW diet, a "biotin-free" version this diet was used in the subsequent streptavidin trial. Neonate *S. lepidus* larvae used in the second and third replicates had had no previous exposure to diets or natural foods containing biotin, but were used directly in the streptavidin trial.

### Trial Designs:

#### *S. lepidus*

35 For the *S. lepidus* streptavidin trial, neonate or 3-day-old larvae were transferred individually to wells of microtitre trays containing "biotin-free" ASW diet. Three replicates were set up, each consisting of 100 larvae receiving a streptavidin treatment and 100 larvae as controls. For the "treatment" group, 0.9 mg/mL streptavidin (55 µM) was blended thoroughly into the diet before it was dispensed into the wells. Control larvae received "biotin-free" ASW diet without any additive. Microtitre trays containing the diets were

first covered by an ironed-on layer of Mylar<sup>®</sup> film. Larvae were then introduced into each well via slits cut in the Mylar<sup>®</sup> and then sealed in by a second covering, this time of Frisk<sup>®</sup> adhesive film. They were observed daily for signs of feeding, burrowing and movement.

- 5 After 15 to 25 days, the films were removed from the trays and each larva was picked out of the diet and placed individually on a small cut block of the same diet in an autoanalyser cup (4 mL). Any deaths were recorded at this time and at approximately weekly intervals thereafter until the end of the experiment (94 days for Replicate 1; 80 days for Replicate 2 and 78 days for Replicate 3).

10

### *L. bonariensis*

- For the *L. bonariensis* streptavidin trial, "biotin-free" ASW diet was made up as before, with the addition of 0.9 mg/mL streptavidin (55  $\mu$ M) for the treatment group. Three replicates, each consisting of 31 streptavidin-fed and 31 control larvae, were set up. For these weevils, 15 each egg was placed directly on a cut block of the appropriate diet in an autoanalyser cup (4 mL) and sealed with its plastic lid. They were examined daily to observe larval hatching, feeding, burrowing or movement. At approximately weekly intervals the containers were opened, the diet block teased apart and larval deaths recorded. Fresh diet of the same type was provided when required. The experiment was ended after 51 days, when 20 many of the control insects were still alive.

### Results:

#### *S. lepidus*

- 25 In each replicate, larval survival was significantly lower for weevils feeding on diet with 55  $\mu$ M streptavidin added than for the control weevils ( $P < 0.001$ , log-rank tests to compare survival curves (Kalbfleisch and Prentice, 1980)). Figure 38 shows the survival curves for all replicates combined. Many of the larvae in the streptavidin treated group appeared to have died during or immediately after a larval moult. Dead larvae often had a soft, 30 transparent head and the darker discarded head capsule attached to the rear of the insect.

- Table 7 shows the median survival times for each group of weevils. Weevils in Replicate 1 had better survival than those in Replicates 2 and 3. This may be due to the Replicate 1 weevils receiving either clover roots or ASW or porina diet with biotin added for three 35 days before the start of the trial. In each case however, larvae treated with streptavidin died significantly sooner than the control larvae.

Control survival was poorer than might be expected for weevils in the field and only four control weevils developed into adults before the end of the experiment, probably because

ASW diet was not the ideal medium for rearing this insect. No adults emerged among the clover root weevils fed streptavidin.

### *L. bonariensis*

- 5 In each replicate, larval survival was significantly lower for the weevils fed streptavidin than for the controls ( $P < 0.001$ , log-rank tests to compare survival curves). Figure 39 shows the survival curves for data from the three replicates combined. As with the clover root weevils, Argentine stem weevil larvae that had received streptavidin appeared to have died during the moulting process and discarded head capsules were found adhering  
10 to the rear ends of dead larvae.

### Conclusions:

- Streptavidin has significant toxicity to the larvae of two plant-eating weevils, the clover root weevil, *S. lepidus*, and the Argentine stem weevil, *L. bonariensis*. This suggests  
15 that pasture plants expressing biotin-binding proteins in the roots or stems could be protected from attack by these pests.

Table 7: Median survival times for *S. lepidus* larvae (days). 95% confidence intervals in brackets

	Replicate 1	Replicate 2	Replicate 3
Streptavidin Treatment	14 (11-15)	4 (4-6)	4 (3-7)
Control	24 (21-29)	8 (4-18)	16 (11-25)

### EXAMPLE 14

Feeding trials with adult clover root weevils (*Sitona lepidus*) (Coleoptera: Curculionidae) fed with avidin-painted clover (*Trifolium repens*) foliage

### Methods:

- 30 Adult *Sitona lepidus* were collected from a field at Ruakura Agricultural Research Centre, Hamilton, New Zealand, using a suction-powered insect-collecting device. They were then placed individually in clear plastic 30 mL containers ("Coulter cups") with vented lids, each containing a single painted leaf of white clover (*Trifolium repens*) with its stem embedded in about 10 mL of 0.4% agar in the bottom of the cup. This kept the leaf turgid  
35 for several days, while providing the weevil with a solid surface to walk on.

The avidin used in this trial was a Calbiochem® product, purchased from Calbiochem-Novabiochem Corporation, La Jolla, CA 92039. It was lyophilized avidin from egg white, Lot 276992.

As the upper surfaces of clover leaves are very hydrophobic, and *S. lepidus* adult weevils typically consume the entire leaf, only the undersides of the leaves were painted. The following solutions were applied with a small sable brush:

- 5        1. Controls were painted with 0.1% (v:v) BondXtra<sup>®</sup> (a wetting, spreading and sticking agent) at a rate of 80 µl solution per g of leaf (fresh weight).
2. "Low" avidin treatment leaves were painted with a 5 mg/mL avidin solution in 0.1 % BondXtra<sup>®</sup> at the same rate as above. This rate approximates a leaf expressing 26 µM avidin.
- 10       3. "High" avidin treatment leaves were painted with a 10 mg/mL avidin solution in 0.1 % BondXtra<sup>®</sup> at the same rate as above. This rate approximates a leaf expressing 52 µM avidin.

15       Between 15 and 18 adult weevils were placed on control leaves. 16 to 18 on low avidin-painted leaves and 16 to 18 on high avidin-painted leaves. The experiment was replicated three times (total of 149 weevils).

Weevils were examined and deaths recorded every weekday until all weevils had died.

## 20       **Results:**

There were no significant differences among the survival curves for adult *S. lepidus* fed clover leaves painted with two doses of avidin or with a control solution without avidin (Figure 40) (log-rank test, Kalbfleisch and Prentice, 1980).

## 25       **Conclusions:**

Avidin is not toxic to adult clover root weevils, *S. lepidus*, when painted onto clover leaves at approximately 26 or 52 µM. It is thus unlikely that transgenic clover plants expressing avidin at these levels will have toxicity to the adult stage of this weevil.

## 30       **EXAMPLE 15**

**Feeding trials with adult honeybees, *Apis mellifera* (Hymenoptera: Apidae), and artificial diet containing avidin**

### **Method:**

- 35       Young adult honeybees were collected as they emerged from frames of capped bee brood taken from hives kept at our apiary in Auckland, New Zealand.

The avidin used in this trial was a Calbiochem<sup>®</sup> product, purchased from Calbiochem-

Novabiochem Corporation, La Jolla, CA 92039. It was lyophilized avidin from egg white, Lot 276992.

5 Bees were assigned randomly to wooden cages (9 x 8 x 7 cm) with mesh on two sides, 30 bees per cage. Each cage was fitted with two gravity feeders, one containing water and the other sugar syrup (60% w:v sucrose solution). These were replenished as necessary during the experiment.

10 Each cage was also provided with a small cup containing a mixture of bee-collected pollen (1 part) and sugar candy (2 parts) (candy recipe: Ambrose, 1992) to which avidin had been added at two different concentrations. One group of cages was supplied with pollen/candy to which 0.1 mg avidin per g of pollen had been added (equivalent to approximately 6.7  $\mu$ M avidin) and a second group was supplied with a mixture containing 0.3 mg avidin per g of pollen (equivalent to approximately 20  $\mu$ M avidin). A third set of bees (controls) 15 received pollen/candy without additive. The trial was replicated four times, i.e. a total of 12 cages of bees.

20 To measure consumption of the pollen/candy food by the bees, each cup was weighed at the start of the experiment and again at Days 8 and 14. Each cage was checked daily for bee deaths.

### Results:

There were no significant differences in the mean quantities of pollen/candy consumed by the three groups of bees (ANOVA) over the first 8 days of exposure to the foods, between 25 Days 8 and 14, or over the entire 14-day period (Figure 41).

Comparisons of survival curves using log-rank tests (Kalbfleisch and Prentice, 1980) showed that bees fed the higher dose of avidin had significantly better survival ( $P < 0.002$ ) than those fed the lower dose. Control bee survival was intermediate between, and did 30 not differ significantly from, that of bees fed either avidin dose (Figure 42).

### Conclusions:

Adult honeybees readily consume pollen/candy mixtures containing approximately 6.7 or 20  $\mu$ M avidin and, when compared with control bees, their survival is unaffected by 35 this consumption. This suggests that if biotin-binding proteins are expressed at these levels in pollen from plants modified to contain these genes, then young adult bees will not be repelled or harmed by such pollen.



## EXAMPLE 16

Feeding trials with slugs (*Deroceras reticulatum*) (Stylommatophora: Agriolimacidae) and snails (*Cantareus aspersus*) (Stylommatophora: Helicidae) fed with avidin painted onto lettuce (*Latuca sativa*) foliage

5

**Methods:**

Snails and slugs were collected from local gardens (Auckland, New Zealand), weighed and placed in groups in sealed plastic containers (220 x 160 x 40 mm) with organically-grown lettuce leaves coated thoroughly with one of the following treatments:

10

1. Controls were painted with 0.1% (v:v) BondXtra® (a wetting, spreading and sticking agent) only;
2. 4.8 µM avidin treatment leaves were painted with an avidin solution in 0.1% BondXtra® that delivered 75 µg of avidin per g fresh weight of lettuce;
- 15 3. 19.2 µM avidin treatment leaves were painted with an avidin solution in 0.1% BondXtra® that delivered 300 µg of avidin per g fresh weight of lettuce.

20

Each container was checked daily for deaths, the interior sprayed with water mist and the painted lettuce replenished as necessary. At the end of the experiment (after 51 days) all surviving animals were weighed.

**Snails:**

- 25 Snails were individually identified with a number written on their shells with permanent marker pen. Two containers of ten snails each were set up for each treatment (i.e. 3 treatments x 2 containers x 10 snails = 60 snails total). Each snail was weighed at the beginning of the experiment and the survivors also weighed at the end.

**Slugs:**

- 30 Five containers, each containing five slugs, were set up for each of the three treatments (i.e. 3 treatments x 5 containers x 5 slugs = 75 slugs total). As slugs could not be individually marked, all five from each container were weighed together at the beginning of the experiment. Surviving slugs were weighed individually at the end of the experiment.

**Results:**

35

**Snails:**

The three groups of snails used in the experiment had similar initial weights (ANOVA, Figure 43). All snails grew during the 51-day experiment and there were no significant

differences in final weights among the three groups (ANOVA, Figure 43).

Few snails died during the experiment (Figure 44). There were no significant differences in mean snail longevity among the three groups (ANOVA, a 51-day longevity was assumed for all snails alive at the end of the experiment, i.e. an underestimate).

### *Slugs:*

Initial weights of slugs were also similar across the three groups, but either lettuce must have been a poor diet for them or the conditions in their containers did not favour their development, because all slugs lost weight during the experiment (Figure 45). There were no significant differences attributable to the treatments in initial or final mean slug weights (ANOVA).

Slug survival, particularly among the controls, was also poor under these experimental conditions (Figure 46). In fact, slugs on lettuce painted with either 4.8  $\mu\text{M}$  or 19.2  $\mu\text{M}$  avidin had significantly greater mean longevity than the control slugs (ANOVA,  $P=0.040$ , assuming all surviving slugs at the end of the experiment had a longevity of 51 days).

### **Conclusions:**

Avidin had no effect on snail growth or survival when applied to their lettuce leaf food at 4.8  $\mu\text{M}$  or 19.2  $\mu\text{M}$  for a period of 51 days.

Slug results were confounded by poor growth of all slugs and poor survival of controls during the trial. However, avidin had no obvious toxicity to these invertebrates over a 51-day period of receiving lettuce painted with 4.8  $\mu\text{M}$  or 19.2  $\mu\text{M}$  of this protein.

**EXAMPLE 17****Evaluation of resistance of tobacco (*Nicotiana tabacum*) plants expressing avidin to three species of root-knot nematodes.****5 Methods:****Plants:**

Tobacco seedlings were germinated either from non-transgenic (NT) seed or from seed collected from three independent selfed original transformant plants (PLA2/1, PLA2/4  
10 and PLA2/24).

**Avidin expression levels:**

Twenty five control and 25 transgenic seedlings were transferred individually to 60-mm-diameter plastic pots of peat based potting mix and left to grow for a week before leaf  
15 samples were taken for ELISA analysis of gene expression (Example 5).

Avidin levels in both roots and leaves were measured earlier in 14 transgenic seedlings (from selfed independent original transformants PLA2/7, PLA2/9 and PLA2/13) and two non-transgenic plants. Levels of avidin varied between 0 and 2.23  $\mu$ M in roots and  
20 0 and 16.84  $\mu$ M in leaves. There was a linear correlation between leaf and root avidin levels in individual plants ( $n = 16$ ,  $R^2 = 0.716$ ). Leaf avidin levels were subsequently used to select experimental material since it is not possible to harvest and measure root material prior to assay. Biotin concentrations in these plants were independent of avidin expression, being 0.05  $\mu$ M in root tissue and 0.7  $\mu$ M in leaves.

25

**Nematodes:**

Twenty highly expressing PLA2 plants and 20 non-transgenic plants were re-potted into 100-mm-diameter pots and a week later inoculated with a suspension 4000 eggs of root-knot nematodes injected into holes around the roots (method described in Sasser and Carter  
30 1985). The nematode species used were *Meloidogyne javanica*, *Meloidogyne hapla* and *Meloidogyne incognita*. Control plants were injected with water. Thus, the design was 3 nematode species + 1 control = 4 inoculation types X 2 gene categories X 5 replicates = 40 pots.

35 After seven weeks, roots were washed free of potting mix and the galls counted. Roots and galls were then crushed with a small roller and extracted in chlorine solution to free the eggs, which were sieved out and counted.

**Results:**

- The levels of avidin in the transgenic plants were  $11.4 \pm 6.8 \mu\text{M}$  (range 2.4 - 26.05). Even at the lowest avidin level, a six-fold molar excess of avidin over biotin can be calculated. There were no significant differences between means of gall and egg counts for each of the three root-knot nematode species ( $P > 0.10$ ) (ANOVA, Sokal and Rohlf 1969) (Table 8). No galls were seen on sham inoculated plants.
- 5

**Conclusion:**

Transgenic tobacco expressing high levels of avidin in root tissue is not resistant to root-knot nematode attack.

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
218

Table 8: Number of eggs laid and galls formed on tobacco roots by three species of nematodes.

<i>M. javanica</i>					<i>M. hapla</i>					<i>M. incognita</i>				
Plant	Galls	Mean (s.e.)	Eggs	Mean (s.e.)	Plant	Galls	Mean	Eggs	Mean (s.e.)	Plant	Galls	Mean (s.e.)	Eggs	Mean
NT					NT					NT				
1	209		0		8	208		87		9	28		0	
2	185		63		10	249		0		15	14		41	
3	127		0		14	256		22		21	22		20	
13	222		0		16	251		0		22	17		34	
17	112	171	56	24	20	247	242	0	22	23	51	26	0	19
		(22)		(15)			(9)		(17)			(7)		(8)
PLA2/					PLA2/					PLA2/				
1/6	139		17		24/9	149		34		1/7	30		16	
4/1	175		42		24/11	216		26		4/8	21		0	
24/1	74		16		24/12	197		65		4/14	22		0	
24/4	89		27		24/14	189		115		24/5	16		25	
24/7	127	121	0	20	4/24	277	206	62	60	24/8	32	24	18	12
		(18)		(7)			(21)		(16)			(3)		(5)

### EXAMPLE 18

Combined toxic effects of avidin expressed in tobacco (*Nicotiana tabacum*) leaves painted with either a protease inhibitor or a Bt insecticidal protein to larval *Helicoverpa armigera* (Lepidoptera: Noctuidae): Bt and avidin act synergistically

#### Constructs:

#### Control lines:

Non-transformed control plants:

Two hundred and forty one plants (NT 201 – NT 441) were grown from seeds produced by selfed NT plant 11 which was used in the trial described in Example 8.

#### Avidin-expressing lines:

Selfed T<sub>2</sub> avidin-expressing generation

Tobacco plants were grown from seeds collected from three of the plants used in the trial described in Example 9. These parent plants were selfed (self-fertilised) and were the T<sub>1</sub> offspring of plants from the original transformant (T<sub>0</sub>) plant lines PLA2/7, PLA2/9 and PLA2/13 used in the trial described in Example 8. The plants used in this trial were thus second-generation (T<sub>2</sub>) selfed plants derived from plants which had been transformed

with the avidin gene with a PPI-I targeting sequence (Example 2).

Ninety eight plants from the PLA2/7 #18 line, 99 from the PLA2/9 #24 line and 126 from the PLA2/13 #22 line were grown for the experiment.

5

On each plant used in the experiment, Leaf 1 was designated as the uppermost (youngest) leaf which was 8cm or more in length from leaf tip to junction of leaf base with the petiole. The leaves below Leaf 1 were assigned numbers consecutively down the plant. Leaves 1, 2 and 3 were used for *H. armigera* feeding, as previous experiments had shown larvae grow best on young leaves.

10

#### Insecticidal proteins:

Two purified insecticidal proteins were painted onto tobacco foliage in this experiment:

15 *Bacillus thuringiensis* insecticidal protein, Cry1Ba. Activated Cry1Ba toxin was obtained from a large-scale fermentation of *B. thuringiensis* Bt4412, purified and cleaved according to the method described by Simpson *et al.* (1997).

Protease inhibitor, aprotinin, obtained from Interger<sup>®</sup> Company, Canada/USA (Product No. 7105, Lot No. NT59808).

20

#### Insects:

*H. armigera* were obtained from a laboratory colony reared on artificial diet as described in McManus and Burgess (1995) and established from moths collected in Christchurch, New Zealand.

25 Neonate *H. armigera* larvae were placed on artificial diet for 48h following emergence from eggs. These late first instar larvae were then placed on tobacco leaves as described below. Initial larval weight was determined as the mean of the individual weights of a randomly chosen sample of 54 of the larvae used in the trial.

#### 30 Determination of avidin expression levels:

Before commencing the experiment, a whole leaf sample comprising a leaf of at least 8cm in length was taken from the 263 of the 323 avidin-expressing plants which had grown the best over an eight week period. Eighty nine PLA2/7 #18 plants, 71 PLA2/9 #24 plants and 103 PLA2/13 #22 plants were tested for avidin expression level using the ELISA assay described in Example 5. The plants were then ranked according to avidin expression level. Plants from the top of the table were then used in treatments requiring "high" expressors and those from the bottom of the table used where "low" expressors were required.

35

**Trial design:**

Larvae were subjected to nine different treatments to test the effects of avidin, aprotinin and Cry1Ba separately and in two-way combinations. Each leaf was weighed before painting, and all solutions were applied at a rate of 100  $\mu$ L solution per g of fresh leaf.

- 5 To ensure leaves remained turgid, the petiole of each cut leaf was immersed in a setting solution of 0.4% w:v agar in a 30 mL coulter cup.

**Treatments:**

- 10 Control tobacco leaves painted with a control solution of 0.1% (v:v) BondXtra<sup>®</sup> (a wetting, spreading and sticking agent).

Control tobacco leaves painted with a 2 mg/ml solution of aprotinin in 0.1% (v:v) BondXtra<sup>®</sup> at the same rate as above. If tobacco leaves are about 2% protein, then this rate approximates a leaf expressing aprotinin as 1% of total soluble protein.

- 15 Control tobacco leaves painted with a 1 mg/ml solution of Cry1Ba in 0.1% (v:v) BondXtra<sup>®</sup> at the same rate as above. If tobacco leaves are about 2% protein, then this rate approximates a leaf expressing Cry1Ba as 0.5% of total soluble protein.

Tobacco leaves expressing avidin at a "low" level (see below) and painted with 0.1% (v:v) BondXtra<sup>®</sup>

- 20 Tobacco leaves expressing avidin at a "low" level and painted with a 2 mg/ml solution of aprotinin in 0.1% (v:v) BondXtra<sup>®</sup>

Tobacco leaves expressing avidin at a "low" level and painted with a 1 mg/ml solution of Cry1Ba in 0.1% (v:v) BondXtra<sup>®</sup>

Tobacco leaves expressing avidin at a "high" level (see below) and painted with 0.1% (v:v) BondXtra<sup>®</sup>

- 25 Tobacco leaves expressing avidin at a "high" level and painted with a 2 mg/ml solution of aprotinin in 0.1% (v:v) BondXtra<sup>®</sup>

Tobacco leaves expressing avidin at a "high" level and painted with a 1 mg/ml solution of Cry1Ba in 0.1% (v:v) BondXtra<sup>®</sup>

- 30 The ranges of avidin expression levels in the plants used were as follows:

	Treatment 4 ("low"):	2.12 – 5.27 $\mu$ M
	Treatment 5 ("low"):	2.62 – 5.30 $\mu$ M
	Treatment 6 ("low"):	3.62 – 5.24 $\mu$ M
35	Treatment 7 ("high"):	12.95 – 21.27 $\mu$ M
	Treatment 8 ("high"):	12.90 – 21.00 $\mu$ M
	Treatment 9 ("high"):	14.18 – 18.10 $\mu$ M

Ten larvae were placed on the underside of each treated leaf inside a 300 x 210 x 80mm plastic storage box lined with paper towels and with a snap-on lid. Three replicate boxes were set up for each treatment, i.e. 27 boxes in total, 30 larvae per treatment (two of the treatments were inadvertently given 31 larvae). Larvae and leaves were checked daily, and leaves were replaced with identically treated fresh leaves from similar plants as necessary so that larvae could feed *ad libitum*.

The experiment was conducted in a controlled temperature room at  $30 \pm 1^\circ\text{C}$  and 60% relative humidity, with a 16:8h light:dark cycle.

10

Larval deaths were recorded on Day 2 and daily thereafter for 14 days or until or all had died if this occurred earlier. Larvae were weighed on Days 3, 6, 8, 10 and 12. Once larvae had begun to pupate in any treatment, larvae in that treatment were no longer weighed.

## 15 Results:

Survival curves for *H. armigera* in the nine different treatment groups are shown in Figure 47. Log-rank tests (Kalbfleisch and Prentice, 1980) were used to compare median survival in the different treatments. The only treatment which did not reduce median survival time compared with control survival was that using aprotinin-painted control leaves.

20

The four treatments using leaves expressing avidin at both high and low levels, with and without aprotinin painted on, killed all larvae within 13 days. Death often occurred during early larval instar moulting. Survival on all these treatments was significantly reduced in comparison with survival on control leaves with and without aprotinin (ANOVA  $P < 0.001$ ) (Payne *et al.* 1993). Median survival times on these four avidin-expressing treatments did not differ significantly from each other. Thus the effect on median larval survival of the combination of avidin expression and aprotinin was equivalent to the effect of avidin expression alone. However, closer examination of the survival curves for the "low avidin" and the "low avidin with aprotinin" reveals that they diverge between days 8 and 12.

25

The proportion of larvae alive on the "low avidin with aprotinin" treatment is significantly lower on days 9, 10 and 11 (ANOVA  $P < 0.05$ ). This demonstrates that avidin can be combined with a protease inhibitor to produce a more toxic effect on larvae, even though the effect of the protease inhibitor alone may be subtle. Additionally, there is no suggestion of antagonism between the two types of resistance protein.

30

The three treatments in which Cry1Ba was painted onto the leaves killed all larvae within four days. Larvae feeding on high and low avidin-expressing leaves painted with Cry1Ba died significantly faster than those feeding on Cry1Ba-painted control leaves (ANOVA  $P < 0.001$ ). The effects on mean larval survival of both the high avidin/Cry1Ba and the low avidin/Cry1Ba combination treatments were greater than the sum of the effects of

40



the high or low avidin expression alone and Cry1Ba painting alone. Thus synergistic effects were observed with these combinations.

5 Growth rates and biomass were plotted for larvae on all the non-Cry1Ba treatments. Larvae feeding on control plants painted with the control solution or the aprotinin solution grew and accumulated biomass exponentially, while those on all treatments expressing avidin at high or low levels failed to grow or accumulate substantial biomass (Figures 48 And 49). Because of the powerful effects of the avidin alone, it was not possible to measure any more subtle effects that the combination with aprotinin may have had on these two  
10 parameters.

### Conclusions:

Synergistic toxic effects on *H. armigera* larvae were observed with combinations of avidin-expressing tobacco leaf and the Bt insecticidal protein. Cry1Ba.

15 This suggests strongly that plants containing chimeric genes and expressing both avidin and Bt will be highly effective in protecting the plants from pest attack. It is likely that such plants will be more toxic than those expressing either protein singly.

20 Avidin at both high and low expression levels when combined with aprotinin was as effective as avidin expression alone in killing larvae and preventing growth and biomass accumulation. During the latter part of the experiment, larval death was greater in the combined aprotinin/low avidin treatment than in the low avidin treatment alone. This demonstrates the possibility that additive or synergistic effects could occur between avidin or streptavidin and a protease  
25 inhibitor which reduces larval growth and survival. The absence of any antagonistic effects between the biotin binding protein and the protease inhibitor shows the compatibility of these two types of resistance factor.

30 It is likely that plants expressing avidin together with a second effective insecticidal protein employing a different mode of action will not only have greater toxicity, but also more durable resistance to pest attack than plants expressing or containing a Bt protein, a protease inhibitor or another type of pest resistance factor on its own.

## REFERENCES

- Argarana C.E., Kuntz, I.D., Birken, Axel R., and Cantor C.R. (1986). Molecular Cloning and Nucleotide Sequence of the Streptavidin Gene Nucleic Acids research 14: 1871-1882.
- Bayer EA, Ben-Hur H, Wilchek M (1990) Isolation and properties of streptavidin. Methods in Enzymology 184: 80-89.
- Bednarek SY, Wilkins TA, Dombrowski JE, Raikhel NV (1990) A carboxyl-terminal propeptide is necessary for proper sorting of barley lectin to vacuoles of tobacco. Plant Cell 1990 2:1145-1155.
- Beuning L. L., Spriggs T. W., and Christeller J. T. (1994) Evolution of the Proteinase Inhibitor I Family and Apparent Lack of Hypervariability in the Proteinase Contact Loop. J. Mol. Evol. 39 (6): 644-654.
- Boulter, D. (1993). Review Article Number 86. Insect Pest Control by Copying Nature using Genetically Engineered Crops. Phytochemistry 34: 1453-1466.
- Broodryk, S.W. 1971. Ecological investigations on the potato tuber moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae). Phytophylactica 3: 73-84.
- Burgess, E.P.J., Main, C.S., Stevens, P.S., Gatehouse, A.M.R., Christeller, J.T. and Laing, W.A. 1993. Protease inhibitors active against porina caterpillar (*Wiseana cervinata*). Proc. 6<sup>th</sup> Australasian Grassl. Invert. Ecol. Conf. 1993 (R.A. Prestidge, Ed.), AgResearch, Hamilton, New Zealand, 331-339.
- Chrispeels, M.J. (1991). Sorting of Proteins in the Secretory System. Annu. Rev. Plant Physiol. Plant Mol. Biol. 42: 21-53.
- Chrispeels MJ, Raihkel NV. Short peptide domains target proteins to plant vacuoles. Cell. 1992 Feb 21; 68(4): 613-616.
- Cleveland, T.E., Thornburg, R.W., and Ryan, C.A. (1987). Molecular Characterization of Wound-inducible Inhibitor I Gene from Potato and the Processing of its mRNA and Protein. Plant Mol. Biol. 8: 199-207.

Dadd, R.H. (1985). Nutrition: Organisms In: Comprehensive Insect Physiology, Biochemistry and Pharmacology (Kerkut G.A. and Gilbert, L.L.) Pergamon Press, NY Vol 4, p313-390).

Ditta G., Stanfield S., Corbin D., and Helinski D.R. (1980) Broad Host Range DNA Cloning System for Gram-Negative Bacteria: Construction of a Gene Bank of *Rhizobium meliloti*. Proc Natl Acad Sci USA 77 (12): 7347-7351.

Donson J, Kearney CM, Hilf ME, Dawson WO (1991); Systematic expression of a bacterial gene by a tobacco mosaic virus-based vector. Proc Natl. Acad. Sci. USA., 88:7204-8.

Drombrowski JE, Raikhel NV. Protein targeting to the plant vacuole – a historical perspective. Braz J. Med. Biol. Res. 1996 Apr; 29(4): 413-430.

Gleave A.P. (1992) A Versatile Binary Vector System with a T-DNA Organisational Structure Conducive to Efficient Integration of Cloned DNA into the Plant Genome. Plant Mol. Biol. 20 (6): 1203-1207.

Gleave, A.P., Mitra, D.S., Markwick, N.P., Morris, B.A.M. and Beuning, L.L. 1998. Enhancing the expression of the *Bacillus thuringiensis* cry9Aa2 gene in transgenic plants by nucleotide sequence modification. Molecular Breeding 4: 459-472.

Gope M. L., Keinänen R. A., Kristo P. A., Conneely O. M., Beattie W. G., Zarucki-Schulz T., O'Malley B. W., and Kulomaa M. S. (1987) Molecular Cloning of the Chicken Avidin cDNA. Nucleic Acids Res. 15 (8): 3595-3606.

Graham, J.S., Pearce, G., Merryweather, J., Titani, K., Ericsson L.H., and Ryan, C.A. (1985). Wound-induced Proteinase Inhibitors from Tomato Leaves. J. Biol. Chem. 260: 6561-6564.

Holwerda BC, Padgett HS, Rogers JC (1992) Proaleurain vacuolar targeting is mediated by short contiguous peptide interactions. Plant Cell 4:307-318.

James, C. and A.F. Krattiger (1996). Global Review of the Field Testing and Commercialisation of Transgenic Plants, 1986 to 1995: The First Decade of Crop Biotechnology. International Service for the Acquisition of Agri-Biotech Applications (ISAAA) Briefs No. 1. 11) ISAAA: Ithaca, NY pp31.

Joshi RL, Joshi V, Ow DW (1990) BSMV genome mediated expression of a foreign gene in dicot and monocot plant cells. EMBO J. 9:2663-9.

Kalbfleisch, J.D. and Prentice, R.L. 1980. Statistical Analysis of Failure Time Data, John Wiley, New York.

Keinanen RA, Wallen MJ, Kristo PA, Laukkanen MO, Toimela TA, Helenius MA, Kulomaa MS Molecular cloning and nucleotide sequence of chicken avidin-related genes. 1-5. Eur.J. Biochem. 220:615-621 (1994).

Keller, M., Sneh, B., Strizhov, N., Prudovsky, E., Regev, A., Koncz, C., Schell, J., Zilberstein, A. (1996): Digestion of delta-endotoxin by gut proteases may explain reduced sensitivity of advanced instar larvae of *Spodoptera littoralis* to CryIC. Insect Biochem. Mol. Biol. 26: 365-73.

Kirsch T, Paris N, Butler JM, Beevers L, Rogers JC. Purification and initial characterization of a potential plant vacuolar targeting receptor. Proc. Natl. Acad. Sci. USA. 1994 Apr 12; 91(8): 3403-3407.

Malone, L.A. and Wigley, P.J. 1990. A practical method for rearing Argentine stem weevil, *Listronotus bonariensis* (Coleoptera: Curculionidae) in the laboratory. *N.Z. Entomol.*, 13, 87-88.

Mariani, C., Goldberg R.B., Leemans, J. (1991), Engineered male sterility in plants. Symp. Soc. Exp. Biol. 45:271-9.

Martilla AT, Airenne KJ, Laitinen OH, Kulik T, Bayer EA, Wilchek M, Kulomaa MS. Engineering of chicken avidin: a progressive series of reduced charge mutants. FEBS Letters 441: 313-317 (1998).

Matsuoka, K., Matsumoto, S., Hattori, T., Machida, Y., and Nakamura, K. (1990) Vacuolar Targeting and Posttranslational Processing of the Precursor to the Sweet Potato Tuberous Root Storage Protein in Heterologous Plant Cells. J. Biol. Chem. 265: 19750-19757.

Meisner, J., K.R.S. Ascher, and D. Lowie. 1974. Phagostimulants for the larva of the potato tuber moth, *Gnorimoschema operculella* Zell. Z. Angew. Entomol. 77: 77-106.

Micheltmore, R., Marsh, E., Seely, S., and Benoit, L. (1987). Transformation of lettuce (*Lactuca sativa*) mediated by *Agrobacterium tumefaciens*. *Plant Cell Reports* 6: 439-442.

Miller, D.W., et al., in *Genetic Engineering* (1986) Setlo W, J K et al., Eds, Plenum Publishing, Vol 8: pages 277-297).

Moore D.S., and Michael S. F. (1995) Mutagenesis of Amplified DNA Sequences Using Ampligase Thermostable DNA Ligase. *Epicentre Forum* 2 (4): 4-5.

Morgan, T.D., B. Oppert, T.H. Czapala, and K.J. Kramer (1993). Avidin and Streptavidin as Insecticidal and Growth Inhibiting Dietary Proteins. *Entomol.exp.appl.* 69: 97-108.

Murray C. and Christeller J. T. (1994) Genomic Nucleotide Sequence of a Proteinase Inhibitor II Gene. *Plant Physiol.* 106: 1681.

Nakamura, K., and Matsuoka, K. (1993) Protein Targeting to the Vacuole in Plant Cells. *Plant Physiol.* 101: 1-5.

Neuhaus JM, Sticher L, Meins F Jr, Boller T (1991) A short C-terminal sequence is necessary and sufficient for the targeting of chitinases to the plant vacuole. *Proc Natl Acad Sci U S A* 88:10362-10366.

Nielsen KJ, Hill JM, Andersom MA, Craik DJ. Synthesis and structure determination of NMR of a putative vacuolar targeting peptide and model of proteinase inhibitor from *Nicotiana glauca*. *Biochemistry*. 1996 Jan 16; 35(2): 369-378.

Rusch SL, Kendall DA. Protein transport via amino-terminal targeting sequences: common themes in diverse systems. *Mol. Membr. Biol.* 1995 Oct; 12(4): 295-307.

Saalebach G, Rosso M, Schumann U (1996) The vacuolar targeting signal of the 2S albumin from Brazil nut resides at the C terminus and involves the C-terminal propeptide as an essential element. *Plant Physiol* 112:975-985.

Sanchez-Serrano, J., Schmidt, R., Schell, J., and Willmitzer, L. (1986). Nucleotide Sequence of Proteinase Inhibitor II Encoding cDNA of Potato (*Solanum tuberosum*) and its Mode of Expression. *Mol. Gen. Genet.* 203: 15-20.

Schatz PJ. (1993) Bio/Technology 11, 1138-1143

Schroder MR, Borkhsenius OW, Matsuoka K, Nakamura K, Raikhel NV. Colocalization of barley lectin and sporamin in vacuoles of transgenic tobacco plants. Plant Physiol. 1993 Feb; 101(2): 451-458.

Seshagiri PB, Adiga PR (1987) Isolation and characterisation of a biotin-binding protein from the pregnant-rat serum and comparison with that from the chicken egg-yolk. Biochim Biophys Acta 916:474-481.

Shao, Z., Cui, Y., Liu, X., Yi, H., Ji, J., Yu, Z. (1998): Processing of delta-endotoxin of *Bacillus thuringiensis* subsp. *Kurstaki* HD-1 in *Heliothis armigera* midgut juice and the effects of protease inhibitors., J Invertebr. Pathol. 72: 73-81.

Subramanian N, Adiga PR (1995) Simultaneous purification of biotin-binding proteins-I and -II from chicken egg yolk and their characterization. Biochem J 308:573-577.

Tague BW, Dickinson CD, Chrispeels MJ (1990) A short domain of the plant vacuolar protein phytohemagglutinin targets invertase to the yeast vacuole. Plant Cell 2:533-546.

Thompson L. D. and Weber P. C. ( 1993) Construction and Expression of a Synthetic Streptavidin-Encoding Gene in *Escherichia coli*. Gene 136: 243-246.

Torres, C., Cantliffe, D.J., Laughner, B., Bieniek, M., Nagata, R., Ashraf, M. and R.J. Ferl (1993). Stable transformation of lettuce cultivar South Bay from cotyledon explants. Plant Cell, Tissue and Organ Culture 34: 279-285.

Turpen, TH. (1999) Tobacco mosaic virus and the virescence of biotechnology. Philos Trans. R. Soc. Lond. Biol. Sci., 354: 665-73, 1999).

Vitale A, Chrispeels MJ (1992) Sorting of proteins to the vacuoles of plant cells. Bioassays 14:151-160.

Von Heijne, G. (1983). Patterns of Amino Acids Near Signal-Sequence Cleavage Sites. Eur. J. Biochem. 133: 17-21.

Walker-Simmons, M., and Ryan, C.A. (1977). Immunological Identification of Proteinase

Inhibitors I and II in Isolated Tomato Leaf Vacuoles. *Plant Physiol.* 60: 61-63.

Wilcheta M, Bayer EA. (eds) 1990. Avidin-Biotin Biotechnology. Methods of Enzymology Vol 184.

Zhang, X., and Conner, A.J. (1992). Genotypic effects on tissue culture response of lettuce cotyledons. J.Genet. and Breed. 46: 287-290.

All references are incorporated herein by reference.